

1. Adams H.A.: Mechanisms of action of ketamine. *Anesth Reanim* 1998; 23: pp. 60-63
[View In Article](#)
2. Admiraal P.V., Knape H., and Zegveld C.: Experience with bezitramide and droperidol in the treatment of severe chronic pain. *Br J Anaesth* 1972 Nov; 44: pp. 1191-1196
[View In Article](#) | [Cross Ref](#)
3. Affaitati G., Fabrizio A., Savini A., et al: A randomized, controlled study comparing a lidocaine patch, a placebo patch, and anesthetic injection for treatment of trigger points in patients with myofascial pain syndrome: evaluation of pain and somatic pain thresholds. *Clin Ther* 2009; 31: pp. 705-720
[View In Article](#) | [Cross Ref](#)
4. Afshar K., Jafari S., Marks A.J., et al: Nonsteroidal anti-inflammatory drugs (NSAIDs) and non-opioids for acute renal colic. *Cochrane Database Syst Rev* 2015 Jun 29; 6:
[View In Article](#)
5. Ahern T., Herring A., Stone M., et al: Effective analgesia with low-dose ketamine and reduced dose hydromorphone in ED patients with severe pain. *Am J Emerg Med* 2013; 31: pp. 847-851
[View In Article](#) | [Cross Ref](#)
6. Ahern T., Herring A., Stone M., et al: Effective analgesia with low-dose ketamine and reduced dose hydromorphone in ED patients with severe pain. *Am J Emerg Med* 2013; 31: pp. 847-851
[View In Article](#) | [Cross Ref](#)
7. Ahern T.L., Herring A.A., Anderson E.S., et al: The first 500: initial experience with widespread use of low-dose ketamine for acute pain management in the ED. *Am J Emerg Med* 2015; 33: pp. 197-201
[View In Article](#) | [Cross Ref](#)
8. Ahern T.L., Herring A.A., Anderson E.S., Madia V.A., Fahimi J., and Fazee B.W.: The First 500: initial experience with widespread use of low-dose ketamine for acute pain management in the ED. *Am J Emerg Med* 2015; 33: pp. 197-201
[View In Article](#) | [Cross Ref](#)
9. Ahern T.L., Herring A.A., Miller S., and Fazee B.W.: Low-dose ketamine infusion for emergency department patients with severe pain. *Pain Med* 2015; 16: pp. 1402-1409
[View In Article](#) | [Cross Ref](#)
10. Ahern T.L., Herring A.A., Miller S., et al: Low-dose ketamine infusion for emergency department patients with severe pain. *Pain Med* 2015; 16: pp. 1402-1409
[View In Article](#) | [Cross Ref](#)
11. Ahern T.L., Herring A.A., Stone M.B., and Fazee B.W.: Effective analgesia with low-dose ketamine and reduced dose hydromorphone in ED patients with severe pain. *Am J Emerg Med* 2013; 31: pp. 847-851
[View In Article](#) | [Cross Ref](#)
12. Ahern T.L., Herring A.A., Stone M.B., and Fazee B.W.: Effective analgesia with low-dose ketamine and reduced dose hydromorphone in ED patients with severe pain. *Am J Emerg Med* 2013; 31: pp. 847-851
[View In Article](#) | [Cross Ref](#)
13. Ahern T.L., Herring A.A., Stone M.B., and Fazee B.W.: Safety and efficacy of low-dose ketamine added to a hydromorphone titration protocol for emergency

department patients with acute severe pain. In (eds): Research forum abstracts . pp. 4
[View In Article](#)

14. Ahern T.L., Herring A.A., Stone M.B., et al: Effective analgesia with low-dose ketamine and reduced dose hydromorphone in ED patients with severe pain. Am J Emerg Med 2013; 31: pp. 847-851
[View In Article](#) | [Cross Ref](#)
15. Ahern TL, Herring AA, Anderson ES, et al. The first 500: initial experience with widespread use of low-dose ketamine for acute pain management in the ED. Am J Emerg Med. 2015;33(2):197–201. [\[PubMed\]](#)
16. Ahern TL, Herring AA, Anderson ES, Madia VA, Fahimi J, Frazee BW. The first 500:initial experience with widespread use of low-dose ketamine for acute pain management in the ED. Am J Emerg Med. 2015;33(2):197–201. [\[PubMed\]](#)
17. Ahern TL, Herring AA, Miller S, et al. Low-dose ketamine infusion for emergency department patients with severe pain. Pain Med. 2015;16(7):1402–9. [\[PubMed\]](#)
18. Ahern TL, Herring AA, Miller S, Frazee BW. Low-dose ketamine infusion for emergency department patients with severe pain. Pain Med. 2015;16(7):1402–9. [\[PubMed\]](#)
19. Ahern TL, Herring AA, Stone M et al. Effective analgesia with low-dose ketamine and reduced dose hydromorphone in ED patients with severe pain. Am J Emerg Med. 2013; 31:847-51.
20. Ahern TL, Herring AA, Stone MB, Frazee BW. Effective analgesia with low-dose ketamine and reduced dose hydromorphone in ED patients with severe pain. *Am J Emerg Med* 2013;31:847-51.
21. Aida S.M., Yamakura T.M., Baba H.M., et al: Preemptive analgesia by intravenous low-dose ketamine and epidural morphine in gastrectomy: a randomized double-blind study. Anesthesiology 2000; 92: pp. 1624-1630
[View In Article](#) | [Cross Ref](#)
22. Akino Y, Teshima T, Kihara A, Kodera-Suzumoto Y, Inaoka M, Higashiyama S, Furusawa Y, Matsuura N (2009) Carbon-ion beam irradiation effectively suppresses migration and invasion of human non-small-cell lung cancer cells. Int J Radiat Oncol Biol Phys 75(2):475–481. <https://doi.org/10.1016/j.ijrobp.2008.12.090> [CrossRef](#)[PubMed](#)[Google Scholar](#)
23. Aksel G, Güler S, Doğan NÖ, Çorbacioğlu SK. A randomized trial comparing intravenous paracetamol, topical lidocaine, and ice application for treatment of pain associated with scorpion stings. Hum Exp Toxicol 2015;34:662-7.
24. Alam A, Gomes T, Zheng H et al. Longterm analgesic use after low-risk surgery: a retrospective cohort study. Arch Intern Med. 172; 2012:425-30.
25. Alam A., Gomes T., Zheng H., et al: Long-term analgesic use after low risk surgery: a retrospective cohort study. Arch Intern Med 2012; 172: pp. 425-430
[View In Article](#) | [Cross Ref](#)
26. Alam A., Gomes T., Zheng H., Mamdani M.M., Juurlink D.N., and Bell C.M.: Long-term analgesic use after low-risk surgery: a retrospective cohort study. Arch Intern Med 2012 Mar 12; 172: pp. 425-430
[View In Article](#) | [Cross Ref](#)

27. Alaska State Board of Nursing advisory opinion. www.commerce.state.ak.us/dnn/Portals/5/pub/NUR_Advisory_Ketamine_Administration.pdf (accessed 2015 May 13)
28. Alexander J, Manno M. Underuse of analgesia in very young pediatric patients with isolated painful injuries. *Ann Emerg Med.* 2003;41:617–622. [PubMed]
29. Alexander R, El-Moalem HE, Gan TJ. Comparison of intravenous ketorolac tromethamine and morphine sulfate in the treatment of postoperative pain. *Pharmacotherapy.* 1990;10:116S–121S. [PubMed]
30. American College of Emergency Physicians Opioid Guideline Writing Panel. Clinical policy: critical issues in the prescribing of opioids for adult patients in the emergency department. *Ann Emerg Med.* 2012; 60:499–525.
31. American College of Emergency Physicians website. 2013 Model of the Clinical Practice of Emergency Medicine. Available at: <http://www.acep.org/Clinical-Practice-Management/Model-of-the-Clinical-Practice-of-Emergency-Medicine>. Accessed November 28, 2015.
[View In Article](#)
32. American College of Emergency Physicians. 2009 Model of the Clinical Practice of Emergency Medicine. Available at:
<http://www.acep.org/WorkArea/linkit.aspx?LinkIdentifier=id&ItemID=47823>. Accessed Feb 23, 2011.
33. American College of Medical Toxicology : Safety issues regarding prescription opioids. *J Med Toxicol* 2016 Mar; 12: pp. 142-144
[View In Article](#) | [Cross Ref](#)
34. American Geriatrics Society Panel on the Pharmacological Management of Persistent Pain in Older Persons : Pharmacological management of persistent pain in older persons. *J Am Geriatr Soc* 2009; 57: pp. 1331-1346
[View In Article](#) | [Cross Ref](#)
35. American Society of Health-System Pharmacists Council on Pharmacy Practice. ASHP statement on pharmacy services to the emergency department. *Am J HealthSyst Pharm.* 2008; 65:2380-3.
36. American Society of Health-System Pharmacists. 2015 ASHP Health-System Pharmacy Initiative. www.ashp.org/s_ashp/cat1c.asp?CID=218&DID=255 (accessed 2008 Jun 6).
37. Amery W.K., Admiraal P.V., Beck P.H., et al: Peroral management of chronic pain by means of bezitramide (R 4845), a long-acting analgesic, and droperidol (R 4749), a neuroleptic. A multicentric pilot-study. *Arzneimittelforschung* 1971 Jun; 21: pp. 868-871
[View In Article](#)
38. Andolfatto G, Willman E, Joo D, et al. Intranasal ketamine for analgesia in the emergency department: a prospective observational series. *Acad Emerg Med* 2013;20:1050–4.
39. Andolfatto G., Willman E., Joo D., et al: Intranasal ketamine for analgesia in the emergency department: a prospective observational series. *Acad Emerg Med* 2013; 20: pp. 1050-1054
[View In Article](#) | [Cross Ref](#)
40. Andolfatto G., Willman E., Joo D., Miller P., Wong W.B., Koehn M., et al: Intranasal ketamine for analgesia in the emergency department: a prospective observational series. *Acad Emerg Med* 2013; 20: pp. 1050-1054
[View In Article](#) | [Cross Ref](#)

41. Argoff C.E., Galer B.S., Jensen M.P., Oleka N., and Gammaitoni A.R.: Effectiveness of the lidocaine patch 5% on pain qualities in three chronic pain states: assessment with the Neuropathic Pain Scale. *Curr Med Res Opin* 2004; 20: pp. S21-S28
[View In Article](#) | [Cross Ref](#)
42. Ault B., and Levy D.: Osteopathic manipulative treatment use in the emergency department: a retrospective medical record review. *J Am Osteopath Assoc* 2015; 115: pp. 132-137
[View In Article](#) | [Cross Ref](#)
43. Averbeck NB, Topsch J, Scholz M, Kraft-Weyrather W, Durante M, Taucher-Scholz G (2016) Efficient rejoining of DNA double-strand breaks despite increased cell-killing effectiveness following spread-out Bragg peak carbon-ion irradiation. *Front Oncol* 6:28. <https://doi.org/10.3389/fonc.2016.00028>
[CrossRef](#)
[PubMed](#)
[PubMed Central](#)
[Google Scholar](#)
44. Azizkhani R, Pourafzali SM, Baloochestani E et al. Comparing the analgesic effect of intravenous acetaminophen and morphine on patients with renal colic pain referring to the emergency department: a randomized controlled trial. *J Res Med Sci*. 2013; 18:772-6.
45. Azizkhani R, Pourafzali SM, Baloochestani E, Masoumi B. Comparing the analgesic effect of intravenous acetaminophen and morphine on patients with renal colic pain referring to the emergency department: a randomized controlled trial. *J Res Med Sci* 2013;18:772-6.
46. Azizkhani R., Pourafzali S.M., Baloochestani E., et al: Comparing the analgesic effect of intravenous acetaminophen and morphine on patients with renal colic pain referring to the emergency department: A randomized controlled trial. *J Res Med Sci* 2013; 18: pp. 772-776
[View In Article](#)
47. Azolosa J.L., Stitzer M.L., and Greenwald M.K.: Opioid physical dependence development: effects of single versus repeated morphine pretreatments and of subjects' opioid exposure history. *Psychopharmacology (Berl)* 1994 Feb; 114: pp. 71-80
[View In Article](#) | [Cross Ref](#)
48. Babu R, Komisarow JM, Agarwal VJ, Rahimpour S, Iyer A, Britt D, Karikari IO, Grossi PM, Thomas S, Friedman AH, Adamson C (2016) Glioblastoma in the elderly: the effect of aggressive and modern therapies on survival. *J Neurosurg* 124(4):998–1007. <https://doi.org/10.3171/2015.4.jns142200>
[CrossRef](#)
[PubMed](#)
[Google Scholar](#)
49. Baehren D.F., Marco C.A., Droz D.E., et al: A statewide prescription monitoring program affects emergency department prescribing behaviors. *Ann Emerg Med* 2010; 56: pp. 19-23
[View In Article](#) | [Cross Ref](#)
50. Baehren D.F., Marco C.A., Droz D.E., et al: A statewide prescription monitoring program affects emergency department prescribing behaviors. *Ann Emerg Med* 2010; 56: pp. 19-23.e1-3
[View In Article](#) | [Cross Ref](#)
51. Bailey B., Bergeron S., Gravel J., et al: Efficacy and impact of intravenous morphine before surgical consultation in children with right lower quadrant pain suggestive of appendicitis: a randomized controlled trial. *Ann Emerg Med* 2007; 50: pp. 371-378
[View In Article](#) | [Cross Ref](#)

52. Bailey J.E., Campagna E., and Dart R.C.: RADARS System Poison Center Investigators. The underrecognized toll of prescription opioid abuse on young children. *Ann Emerg Med* 2009 Apr; 53: pp. 419-424
[View In Article](#) | [Cross Ref](#)
53. Baker H.: Lotus Illustrated Medical Dictionary. New Delhi, India: Lotus Press, 2006.
[View In Article](#)
54. Baley K, Michalov K, Kossick MA, McDowell M. Intravenous acetaminophen and intravenous ketorolac for management of pediatric surgical pain: a literature review. *AANA J* 2014;82:53-64.
55. Balshem H, Helfand M, Schünemann HJ, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011;64:401-6.
56. Balshem H., Helfand M., Schunemann H.J., et al: GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011; 64: pp. 401-406
[View In Article](#) | [Cross Ref](#)
57. Balshem H., Helfand M., Schünemann H.J., et al: GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011; 64: pp. 401-406
[View In Article](#) | [Cross Ref](#)
58. Baranowski A.P., De Courcey J., and Bonello E.: A trial of intravenous lidocaine on the pain and allodynia of postherpetic neuralgia. *J Pain Symptom Manage* 1999; 17: pp. 429-433
[View In Article](#) | [Cross Ref](#)
59. Barazzuol L, Jena R, Burnet NG, Jeynes JC, Merchant MJ, Kirkby KJ, Kirkby NF (2012) In vitro evaluation of combined temozolomide and radiotherapy using X rays and high-linear energy transfer radiation for glioblastoma. *Radiat Res* 177(5):651–662[CrossRef](#)[PubMed](#)[Google Scholar](#)
60. Barnett M.L., Olenski A.R., and Jena A.B.: Opioid-prescribing patterns of emergency physicians and risk of long-term use. *N Engl J Med* 2017; 376: pp. 663-673
[View In Article](#) | [Cross Ref](#)
61. Bates C., Laciak R., Southwick A., and Bishoff J.: Overprescription of postoperative narcotics: a look at postoperative pain medication delivery, consumption and disposal in urological practice. *J Urol* 2011 Feb; 185: pp. 551-555
[View In Article](#) | [Cross Ref](#)
62. Baulch JE, Geidzinski E, Tran KK, Yu L, Zhou YH, Limoli CL (2016) Irradiation of primary human gliomas triggers dynamic and aggressive survival responses involving microvesicle signaling. *Environ Mol Mutagenesis* 57(5):405–415. <https://doi.org/10.1002/em.21988> [CrossRef](#)[Google Scholar](#)
63. Baumann BM, Holmes JH, Chansky ME, Levey H, Kulkarni M, Boudreaux ED. Pain assessments and the provision of analgesia: the effects of a templated chart. *Acad Emerg Med*. 2007;14:47–52. [\[PubMed\]](#)
64. Bean-Lijewski JD, Hunt RD. Effect of ketorolac on bleeding time and postoperative pain in children:a double-blind, placebo-controlled comparison with meperidine. *J Clin Anesth.* 1996;8(1):25–30. [\[PubMed\]](#)
65. Beauchamp GA, Winstanley EL, Ryan SA, Lyons MS. Moving beyond misuse and diversion: the urgent need to consider the role of iatrogenic addiction in the current opioid epidemic. *Am J Public Health*. 2014; 104:2023-7.

66. Beaudoin F., Lin C., Guan W., et al: Low-dose ketamine improves pain relief in patients receiving intravenous opioids for acute pain in the emergency department: results of a randomized, double-blind, clinical trial. *Soc Acad Emerg Med* 2014; 21: pp. 1193-1202
[View In Article](#) | [Cross Ref](#)
67. Beaudoin F.L., Lin C., Guan W., and Merchant R.C.: Low-dose ketamine improves pain relief in patients receiving intravenous opioids for acute pain in the emergency department: results of a randomized, double-blind, clinical trial. *Acad Emerg Med* 2014; 21: pp. 1193-1202
[View In Article](#) | [Cross Ref](#)
68. Beaudoin F.L., Lin C., Guan W., and Merchant R.C.: Low-dose ketamine improves pain relief in patients receiving intravenous opioids for acute pain in the emergency department: results of a randomized, double-blind, clinical trial. *Acad Emerg Med* 2014; 21: pp. 1193-1202
[View In Article](#) | [Cross Ref](#)
69. Beaudoin F.L., Lin C., Guan W., et al: Low-dose ketamine improves pain relief in patients receiving intravenous opioids for acute pain in the emergency department: results of a randomized, double-blind, clinical trial. *Acad Emerg Med* 2014; 21: pp. 1193-1202
[View In Article](#) | [Cross Ref](#)
70. Beaudoin F.L., Lin C., Guan W., et al: Low-dose ketamine improves pain relief in patients receiving intravenous opioids for acute pain in the emergency department: results of a randomized, double-blind, clinical trial. *Acad Emerg Med* 2014; 21: pp. 1193-1202
[View In Article](#) | [Cross Ref](#)
71. Beaudoin F.L., Lin C., Guan W., et al: Low-dose ketamine improves pain relief in patients receiving intravenous opioids for acute pain in the emergency department: results of a randomized, double-blind, clinical trial. *Acad Emerg Med* 2014; 21: pp. 1193-1202
[View In Article](#) | [Cross Ref](#)
72. Beaudoin F.L., Nagdev A., Merchant R.C., et al: Ultrasound-guided femoral nerve blocks in elderly patients with hip fractures. *Am J Emerg Med* 2010; 28: pp. 76-81
[View In Article](#) | [Cross Ref](#)
73. Beaudoin FL, Lin C, Guan W, et al. Low-dose ketamine improves pain relief in patients receiving intravenous opioids for acute pain in the emergency department: results of a randomized, double-blind, clinical trial. *Acad Emerg Med.* 2014;21(11):1193–202.
[PubMed]
74. Beaudoin FL, Lin C, Guan W, Merchant RC. Low-dose ketamine improves pain relief in patients receiving intravenous opioids for acute pain in the emergency department: results of a randomized, double-blind, clinical trial. *Acad Emerg Med.* 2014;21(11):1193–202.
[PubMed]
75. Beel TL, Mitchiner JC, Frederiksen SM, McCormick J. Patient preferences regarding pain medication in the ED. *Am J Emerg Med.* 2000;18:376–380. [PubMed]
76. Beeson M.S., Carter W.A., Christopher T.A., et al: Emergency medicine milestones. *J Grad Med Educ* 2013; 5: pp. 5-13
[View In Article](#) | [Cross Ref](#)
77. Bektas F, Eken C, Karadeniz O, Goksu E, Cubuk M, Cete Y. Intravenous paracetamol or morphine for the treatment of renal colic: a randomized, placebo-controlled trial. *Ann Emerg Med* 2009;54:568–74.

78. Bektas F, Eken C, Karedeniz O et al. Intravenous paracetamol or morphine for the treatment of renal colic: a randomized, placebo-controlled trial. Ann Emerg Med. 2009; 54:569-74.
79. Bektas F., Eken C., Karadeniz O., et al: Intravenous paracetamol or morphine for the treatment of renal colic: a randomized, placebo-controlled trial. Ann Emerg Med 2009; 54: pp. 568-574
[View In Article](#) | [Cross Ref](#)
80. Bell R., Montoya D., Shuaib A., et al: A comparative trial of three agents in the treatment of acute migraine headache. Ann Emerg Med 1990; 19: pp. 1079-1082
[View In Article](#) | [Cross Ref](#)
81. Bell R.F., Dahl J.B., Moore R.A., and Kalso E.: Perioperative ketamine for acute postoperative pain. Cochrane Database Syst Rev 2009; undefined: pp. CD004603
[View In Article](#)
82. Bennett GJ. Update on the neurophysiology of pain transmission and modulation: focus on the NMDA-receptor. *J Pain Symptom Manage* 2000;19:S2-S6.
83. Benzina S, Altmeyer A, Malek F, Dufour P, Denis J-M, Guelette J, Bischoff P (2008) High-LET radiation combined with oxaliplatin induce autophagy in U-87 glioblastoma cells. *Cancer Lett* 264(1):63–70. <https://doi.org/10.1016/j.canlet.2008.01.023> CrossRefPubMedGoogle Scholar
84. Berland D., and Rodgers P.: Rational use of opioids for management of chronic non-terminal pain. *Am Fam Physician* 2012; 86: pp. 252-258
[View In Article](#) | [Cross Ref](#)
85. Berman 1990.
[View In Article](#)
86. Berman W., Fripp R.R., Rubler M., and Alderete L.: Hemodynamic effects of ketamine in children undergoing cardiac catheterization. *Pediatr Cardiol* 1990; 11: pp. 72-76
[View In Article](#) | [Cross Ref](#)
87. Berry NS, Folstad JE, Bauman JL et al. Follow-up observations on 24-hour pharmacotherapy services in the emergency department. *Ann Pharmacother*. 1992; 26:476-80.
88. Bhakta HC, Marco CA. Pain management: association with patient satisfaction among emergency department patients. *J Emerg Med*. 2014; 46:456-64.
89. Bhoi S., Sinha T.P., Rodha M., et al: Feasibility and safety of ultrasound-guided nerve block for management of limb injuries by emergency care physicians. *J Emerg Trauma Shock* 2012; 5: pp. 28-32
[View In Article](#) | [Cross Ref](#)
90. Biering K., Hjollund N.H., and Frydenberg M.: Using multiple imputation to deal with missing data and attrition in longitudinal studies with repeated measures of patient-reported outcomes. *Clin Epidemiol* 2015; 7: pp. 91-106
[View In Article](#) | [Cross Ref](#)
91. Bijur P, Bérard A, Esses D, Calderon Y, Gallagher EJ. Race, ethnicity, and management of pain from long-bone fractures: A prospective study of two academic urban emergency departments. *Acad Emerg Med*. 2008;15:589–597. [PubMed]
92. Bijur P.E., Kenny M.K., and Gallagher E.J.: Intravenous morphine at 0.1 mg/kg is not effective for controlling severe acute pain in the majority of patients. *Ann Emerg Med*

2005; 46: pp. 362-367

[View In Article](#) | [Cross Ref](#)

93. Bijur P.E., Kenny M.K., and Gallagher E.J.: Intravenous morphine at 0.1mg/kg is not effective for controlling acute pain in the majority of patients. Ann Emerg Med 2005; 46: pp. 362-365

[View In Article](#) | [Cross Ref](#)

94. Bijur P.E., Kenny M.K., and Gallagher E.J.: Intravenous morphine at 0.1 mg/kg is not effective for controlling severe acute pain in the majority of patients. Ann Emerg Med 2005; 46: pp. 362-367

[View In Article](#) | [Cross Ref](#)

95. Bijur P.E., Latimer C.T., and Gallagher E.J.: Validation of a verbally administered numerical rating scale of acute pain for use in the emergency department. Acad Emerg Med 2003; 10: pp. 390-392

[View In Article](#)

96. Bijur P.E., Latimer C.T., and Gallagher E.J.: Validation of a verbally administered numerical rating scale of acute pain for use in the emergency department. Acad Emerg Med 2003; 10: pp. 390-392

[View In Article](#)

97. Bijur P.E.: Validation of a verbally administered numerical rating scale of acute pain for use in the emergency department. Acad Emerg Med 2003; 10: pp. 390-392

[View In Article](#)

98. Bijur PE, Kenny MK, Gallagher EJ. Intravenous morphine at 0.1 mg/kg is not effective for controlling severe acute pain in the majority of patients. Ann Emerg Med. 2005;46:362–367. [\[PubMed\]](#)

99. Birnbaum A., Esses D., Bijur P.E., et al: Randomized double-blind placebo- controlled trial of two intravenous morphine dosages (0.10 mg/kg and 0.15 mg/kg) in emergency department patients with moderate to severe acute pain. Ann Emerg Med 2007; 49: pp. 445-453

[View In Article](#) | [Cross Ref](#)

100. Birnbaum A., Esses D., Bijur P.E., et al: Randomized double-blind placebo- controlled trial of two intravenous morphine dosages (0.10 mg/kg and 0.15 mg/kg) in emergency department patients with moderate to severe acute pain. Ann Emerg Med 2007; 49: pp. 445-453

[View In Article](#) | [Cross Ref](#)

101. Birnbaumer D.M.: IV Ketamine is effective for procedural sedation in adults. Emerg Med J 2008; undefined: pp. 26

[View In Article](#)

102. Blaivas M., and Lyon M.: Ultrasound-guided interscalene block for shoulder dislocation reduction in the ED. Am J Emerg Med 2006; 24: pp. 293-296

[View In Article](#) | [Cross Ref](#)

103. Blank FS, Mader TJ, Wolfe J, Keyes M, Kirschner R, Provost D. Adequacy of pain assessment and pain relief and correlation of patient satisfaction in 68 ED fast-track patients. J Emerg Nurs. 2001;27:327–334. [\[PubMed\]](#)

104. Bogenschutz M.P., Donovan D.M., Mandler R.N., et al: Brief intervention for patients with problematic drug use presenting in emergency departments: a randomized

clinical trial. JAMA Intern Med 2014; 174: pp. 1736-1745

[View In Article](#) | [Cross Ref](#)

105. Bond CA, Raehl CL, Pitterle ME. Staffing and the cost of clinical and hospital pharmacy services in United States hospitals. Pharmacotherapy. 1999; 19:767-81.
106. Bond G.R., Ho M., and Woodward R.W.: Trends in hepatic injury associated with unintentional overdose of paracetamol (Acetaminophen) in products with and without opioid: an analysis using the National Poison Data System of the American Association of Poison Control Centers, 2000–7. Drug Saf. 2012 Feb 1; 35: pp. 149-157
[View In Article](#) | [Cross Ref](#)
107. Borland M., Jacobs I., King B., and O'Brien D.: A randomized controlled trial comparing intranasal fentanyl to intravenous morphine for managing acute pain in children in the emergency department. Ann Emerg Med 2007; 49: pp. 335-340
[View In Article](#) | [Cross Ref](#)
108. Borland M., Jacobs I., King B., and O'Brien D.: A randomized controlled trial comparing intranasal fentanyl to IV morphine for managing acute pain in children in the emergency department. Ann Emerg Med 2007; 49: pp. 335-340
[View In Article](#) | [Cross Ref](#)
109. Borland M., Jacobs I., King B., et al: A randomized controlled trial comparing intranasal fentanyl to intravenous morphine for managing acute pain in children in the emergency department. Ann Emerg Med 2007; 49: pp. 335-340
[View In Article](#) | [Cross Ref](#)
110. Borland M., Jacobs I., King B., et al: A randomized controlled trial comparing intranasal fentanyl to intravenous morphine for managing acute pain in children in the emergency department. Ann Emerg Med 2007; 49: pp. 335-340
[View In Article](#) | [Cross Ref](#)
111. Boutureira J., Trim C.M., and Cornell K.K.: Acute pulmonary edema after diazepam-ketamine in a dog. Vet Anesth Analg 2006; 34: pp. 371-376
[View In Article](#)
112. Bovill J.G.: Intravenous anesthesia for the patient with left ventricular dysfunction. Semin Cardiothorac Vasc Anesth. 2006; 10: pp. 43-48
[View In Article](#) | [Cross Ref](#)
113. Bowers KJ, McAllister KB, Ray M, Heitz C. Ketamine as an adjunct to opioids for acute pain in the emergency department:a randomized controlled trial. Acad Emerg Med. 2017;24(6):676–85. [\[PubMed\]](#)
114. Boyer E.W.: Management of opioid analgesic overdose. N Engl J Med 2012 Jul 12; 367: pp. 146-155
[View In Article](#) | [Cross Ref](#)
115. Brady J.E., DiMaggio C.J., Keyes K.M., Doyle J.J., Richardson L.D., and Li G.: Emergency department utilization and subsequent prescription drug overdose death. Ann Epidemiol 2015 Aug; 25: pp. 613-619
[View In Article](#) | [Cross Ref](#)
116. Brands B., Paglia-Boak A., Sproule B.A., Leslie K., and Adlaf E.M.: Nonmedical use of opioid analgesics among Ontario students. Can Fam Physician 2010 Mar; 56: pp. 256-262
[View In Article](#)

117. Bredlau A, McDermott MP, Adams HR, et al. Oral ketamine for children with chronic pain: a pilot phase 1 study. *J Pediatr* 2013;163:194–200.
118. Brennan M.J., and Stanos S.: Strategies to optimize pain management with opioids while minimizing risk of abuse. *PM R* 2010 Jun; 2: pp. 544-558
[View In Article](#) | [Cross Ref](#)
119. Brown C.R., Moodie J.E., Wild V.M., et al: Comparison of intravenous ketorolac tromethamine and morphine sulfate in the treatment of postoperative pain. *Pharmacotherapy* 1990; 10: pp. 116S-121S
[View In Article](#)
120. Brown JN, Barnes CL, Beasley B et al. Effect of pharmacists on medication errors in an emergency department. *Am J Health-Syst Pharm*. 2008; 65:330-3.
121. Brozek JL, Akl EA, Alonso-Coello P, et al. Grading quality of evidence and strength of recommendations in clinical practice guidelines. Part 1 of 3. An overview of the GRADE approach and grading quality of evidence about interventions. *Allergy* 2009;64:669-77.
122. Brush D.E.: Complications of long-term opioid therapy for management of chronic pain: the paradox of opioid-induced hyperalgesia. *J Med Toxicol* 2012 Dec; 8: pp. 387-392
[View In Article](#) | [Cross Ref](#)
123. Burghardt L.C., Ayers J.W., Brownstein J.S., Bronstein A.C., Ewald M.B., and Bourgeois F.T.: Adult prescription drug use and pediatric medication exposures and poisonings. *Pediatrics* 2013 Jul; 132: pp. 18-27
[View In Article](#) | [Cross Ref](#)
124. Butler M.M., Ancona R.M., Beauchamp G.A., Yamin C.K., Winstanley E.L., Hart K.W., et al: Emergency department prescription opioids as an initial exposure preceding addiction. *Ann Emerg Med* 2016 Feb 10; undefined:
[View In Article](#)
125. Calcaterra S.L., Yamashita T.E., Min S.J., Keniston A., Frank J.W., and Binswanger I.A.: Opioid prescribing at hospital discharge contributes to chronic opioid use. *J Gen Intern Med* 2016 May; 31: pp. 478-485
[View In Article](#) | [Cross Ref](#)
126. Canfield M.C., Keller C.E., Frydrych L.M., Ashrafioun L., Purdy C.H., and Blondell R.D.: Prescription opioid use among patients seeking treatment for opioid dependence. *J Addict Med* 2010 Jun; 4: pp. 108-113
[View In Article](#) | [Cross Ref](#)
127. Cantrill S.V., Brown M.D., Carlisle R.J., et al: Clinical policy: critical issues in the prescribing of opioids for adult patients in the emergency department. *Ann Emerg Med* 2012; 60: pp. 499-525
[View In Article](#) | [Cross Ref](#)
128. Carroll I, Barelka P, Meng Want CK et al. A pilot cohort study of the determinants of longitudinal opioid use after surgery. *Anesth Analg*. 115:694-702.
129. Carroll I., Barelka P., Wang C.K., Wang B.M., Gillespie M.J., McCue R., et al: A pilot cohort study of the determinants of longitudinal opioid use after surgery. *Anesth Analg* 2012 Sep; 115: pp. 694-702
[View In Article](#)

130. Carter MK, Allin DM, Scott LA et al. Pharmacist-acquired medication histories in a university hospital emergency department. *Am J Health-Syst Pharm.* 2006; 63:2500-3.
131. Castellsague J., Riera-Guardia N., Calingaert B., et al: Individual NSAIDs and upper gastrointestinal complications: a systematic review and meta-analysis of observational studies (the SOS project). *Drug Saf* 2012; 35: pp. 1127-1146
[View In Article](#)
132. Castellsague J., Riera-Guardia N., Calingaert B., et al: Individual NSAIDs and upper gastrointestinal complications: a systematic review and meta-analysis of observational studies (the SOS project). *Drug Saf* 2012; 35: pp. 1127-1146
[View In Article](#)
133. Castellsague J., Riera-Guardia N., Calingaert B., et al: Individual NSAIDs and upper gastrointestinal complications: a systematic review and meta-analysis of observational studies (the SOS project). *Drug Saf* 2012; 35: pp. 1127-1146
[View In Article](#)
134. Catan T., and Perez E.: A pain-drug champion has second thoughts. *Wall Str. J* 2012; 15:
[View In Article](#)
135. Catapano M.S.: The analgesic efficacy of ketorolac for acute pain [review]. *J Emerg Med* 1996; 14: pp. 67-75
[View In Article](#) | [Cross Ref](#)
136. Catapano M.S.: The analgesic efficacy of ketorolac for acute pain. *J Emerg Med* 1996; 14: pp. 67-75
[View In Article](#) | [Cross Ref](#)
137. Catapano M.S.: The analgesic efficacy of ketorolac for acute pain. *J Emerg Med* 1996; 14: pp. 67-75
[View In Article](#) | [Cross Ref](#)
138. Catapano MS. The analgesic efficacy of Ketorolac for acute pain. *J Emerg Med.* 1996;14(1):67-75. [[PubMed](#)]
139. Cavalieri TA. Management of pain in older adults. *J Am Osteopath Assoc.* 2005;105(3 Suppl 1):S12. [[PubMed](#)]
140. Cawthorn T.R., Phelan R., Davidson J.S., et al: Retrospective analysis of perioperative ketorolac and postoperative bleeding in reduction mammoplasty. *Can J Anaesth* 2012; 59: pp. 466-472
[View In Article](#) | [Cross Ref](#)
141. Cawthorn TR, Phelan R, Davidson JS, Turner KE. Retrospective analysis of perioperative ketorolac and postoperative bleeding in reduction mammoplasty. *Can J Anaesth.* 2012;59(5):466-72. [[PubMed](#)]
142. Center for Disease Control and Prevention : Opioid Overdose Fact Sheet.
<http://www.cdc.gov/drugoverdose/index.html>
[View In Article](#)
143. Centers for Disease Control and Prevention (CDC) : CDC guideline for prescribing opioids for chronic pain.
<https://www.cdc.gov/drugoverdose/prescribing/guideline.html>
[View In Article](#)
144. Centers for Disease Control and Prevention website. National Hospital Ambulatory Medical Care Survey 2011. Available at:
<http://www.cdc.gov/nchs/fastats/emergency-department.htm>. Accessed November

28, 2015.

[View In Article](#)

145. Centers for Disease Control and Prevention website. Prescription drug overdose. Available at: <http://www.cdc.gov/drugoverdose/index.html>. Accessed November 28, 2015.
[View In Article](#)
146. Centers for Disease Control and Prevention. National Ambulatory Medical Care Survey: 2012 state and national summary tables. www.cdc.gov/nchs/ahcd.htm (accessed 2014 Nov 1).
147. Centers for Disease Control and Prevention. Public Health Law: doctor shopping laws. www.cdc.gov/phlp/docs/menushoppinglaws.pdf (accessed 2015 Apr 28).
148. Cepeda MS, Carr DB, Miranda N et al. Comparison of morphine, ketorolac, and their combination for postoperative pain. *Anesthesiology*. 2005; 103:1225-32.
149. Challapalli V., Tremont-Lukats I.W., McNicol E.D., et al: Systemic administration of local anesthetic agents to relieve neuropathic pain. *Cochrane Database Syst Rev* 2005; undefined:
[View In Article](#)
150. Chan L, Verdile VP. Do patients receive adequate pain control after discharge from the ED? *Am J Emerg Med* 1998; 16:705-7.
151. Chang A.K., Bijur P.E., Campbell C.M., et al: Safety and efficacy of rapid titration using 1mg doses of intravenous hydromorphone in emergency department patients with acute severe pain: the “1+1” protocol. *Ann Emerg Med* 2009; 54: pp. 221-225
[View In Article](#) | [Cross Ref](#)
152. Chang A.K., Bijur P.E., Napolitano A., Lupow J., et al: Two milligrams i.v. hydromorphone is efficacious for treating pain but is associated with oxygen desaturation. *J Opioid Manag* 2009; 5: pp. 75-80
[View In Article](#) | [Cross Ref](#)
153. Chang H.Y., Daubresse M., Kruszewski S.P., and Alexander G.C.: Prevalence and treatment of pain in EDs in the United States, 2000 to 2010. *Am J Emerg Med* 2014 May; 32: pp. 421-431
[View In Article](#) | [Cross Ref](#)
154. Chang H.Y., Daubresse M., Kruszewski S.P., et al: Prevalence and treatment of pain in EDs in the United States, 2000 to 2010. *Am J Emerg Med* 2014; 32: pp. 421-431
[View In Article](#) | [Cross Ref](#)
155. Chang H.Y., Daubresse M., Kruszewski S.P., et al: Prevalence and treatment of pain in EDs in the United States, 2000 to 2010. *Am J Emerg Med* 2014; 32: pp. 421-431
[View In Article](#) | [Cross Ref](#)
156. Cheng D., and Majlesi N.: Emergency department opioid prescribing guidelines for the treatment of non-cancer related pain. Milwaukee (WI): AAEM Position Statement, 2013.
[View In Article](#)
157. Chisholm CD, Collison EK, Nelson DR, Cordell WH. Emergency department workplace interruptions: are emergency physicians “interrupt-driven” and “multitasking”? *Acad Emerg Med*. 2000;7:1239–1243. [[PubMed](#)]

158. Chizh BA, Headley PM. NMDA antagonists and neuropathic pain-multiple drug targets and multiple uses. *Curr Pharm Des* 2005;11:2977–94.
159. Chou R., Turner J.A., Devine E.B., Hansen R.N., Sullivan S.D., Blazina I., et al: The effectiveness and risks of long-term opioid therapy for chronic pain: a systematic review for a National Institutes of Health Pathways to Prevention Workshop. *Ann Intern Med* 2015 Feb 17; 162: pp. 276-286
[View In Article](#) | [Cross Ref](#)
160. Chudnofsky C.R., Weber J.E., Stoyanoff P.J., Colone P.D., Wilkerson M.D., Hallinen D.L., et al: A combination of midazolam and ketamine for procedural sedation and analgesia in adult emergency department patients. *Acad Emerg Med* 2000; 7: pp. 228-235
[View In Article](#) | [Cross Ref](#)
161. Cicero T.J., Ellis M.S., Surratt H.L., and Kurtz S.P.: Factors influencing the selection of hydrocodone and oxycodone as primary opioids in substance abusers seeking treatment in the United States. *Pain* 2013 Dec; 154: pp. 2639-2648
[View In Article](#) | [Cross Ref](#)
162. Clarke H., Soneji N., Ko D.T., Yun L., and Wijeysundera D.N.: Rates and risk factors for prolonged opioid use after major surgery: population based cohort study. *BMJ* 2014 Feb 11; 348: pp. g1251
[View In Article](#) | [Cross Ref](#)
163. Combs SE, Bohl J, Elsasser T, Weber KJ, Schulz-Ertner D, Debus J, Weyrather WK (2009) Radiobiological evaluation and correlation with the local effect model (LEM) of carbon ion radiation therapy and temozolomide in glioblastoma cell lines. *Int J Radiat Biol* 85(2):126–
137. <https://doi.org/10.1080/09553000802641151> [CrossRef](#)[PubMed](#)[Google Scholar](#)
164. Combs SE, Bruckner T, Mizoe JE, Kamada T, Tsujii H, Kieser M, Debus J (2013) Comparison of carbon ion radiotherapy to photon radiation alone or in combination with temozolomide in patients with high-grade gliomas: explorative hypothesis-generating retrospective analysis. *Radiother Oncol* 108(1):132–135. <https://doi.org/10.1016/j.radonc.2013.06.026> [CrossRef](#)[PubMed](#)[Google Scholar](#)
165. Combs SE, Kessel K, Habermehl D, Haberer T, Jakel O, Debus J (2013) Proton and carbon ion radiotherapy for primary brain tumors and tumors of the skull base. *Acta Oncol* 52(7):1504–1509. <https://doi.org/10.3109/0284186x.2013.818255> [CrossRef](#)[PubMed](#)[Google Scholar](#)
166. Combs SE, Kieser M, Rieken S, Habermehl D, Jakel O, Haberer T, Nikoghosyan A, Haselmann R, Unterberg A, Wick W, Debus J (2010) Randomized phase II study evaluating a carbon ion boost applied after combined radiochemotherapy with temozolomide versus a proton boost after radiochemotherapy with temozolomide in patients with primary glioblastoma: the CLEOPATRA trial. *BMC Cancer* 10:478. <https://doi.org/10.1186/1471-2407-10-478> [CrossRef](#)[PubMed](#)[MedPub](#)[Central](#)[Google Scholar](#)
167. Compton P., Miotti K., and Elashoff D.: Precipitated opioid withdrawal across acute physical dependence induction methods. *Pharmacol Biochem Behav* 2004 Feb; 77:

pp. 263-268

[View In Article](#) | [Cross Ref](#)

168. CONSORT : <http://www.consort-statement.org/consort-2010>
[View In Article](#)
169. Cordell W.H., Keene K.K., Giles B.K., et al: The high prevalence of pain in emergency medical care. *Am J Emerg Med* 2012; 20: pp. 165-169
[View In Article](#) | [Cross Ref](#)
170. Cordell W.H., Keene K.K., Giles B.K., Jones J.B., Jones J.H., and Brizendine E.J.: The high prevalence of pain in emergency medical care. *Am J Emerg Med* 2002; 20: pp. 165-169
[View In Article](#) | [Cross Ref](#)
171. Cordell W.H., Keene K.K., Giles B.K., Jones J.B., Jones J.H., and Brizendine E.J.: The high prevalence of pain in emergency medical care. *Am J Emerg Med* 2002; 20: pp. 165-169
[View In Article](#) | [Cross Ref](#)
172. Cordell WH, Keene KK, Giles BK, Jones JB, Jones JH, Brizendine Ej. The high prevalence of pain in emergency medical care. *Am J Emerg Med*. 2002; 20:165-9.
173. Cordes N, Hansmeier B, Beinke C, Meineke V, van Beuningen D (2003) Irradiation differentially affects substratum-dependent survival, adhesion, and invasion of glioblastoma cell lines. *Br J Cancer* 89(11):2122–2132. <https://doi.org/10.1038/sj.bjc.6601429>
[CrossRef](#)
[PubMed](#)
[PubMed Central](#)
[Google Scholar](#)
174. Corrigan M., Wilson S.S., and Hampton J.: Safety and efficacy of intranasally administered medications in the emergency department and prehospital settings. *Am J Health Syst Pharm* 2015; 72: pp. 1544-1554
[View In Article](#) | [Cross Ref](#)
175. Craig M, Jeavons R, Probert J et al. Randomised comparison of intravenous paracetamol and intravenous morphine for acute traumatic limb pain in the emergency department. *Emerg Med J*. 2012; 29:37-9.
176. Craig M, Jeavons R, Probert J, Benger J. Randomised comparison of intravenous paracetamol and intravenous morphine for acute traumatic limb pain in the emergency department. *Emerg Med J* 2012;29:37-9.
177. Craig M., Jeavons R., Probert J., and Benger J.: Randomised comparison of intravenous paracetamol and intravenous morphine for acute traumatic limb pain in the emergency department. *Emerg Med J* 2012 Jan; 29: pp. 37-39
[View In Article](#) | [Cross Ref](#)
178. Craig M., Jeavons R., Probert J., et al: Randomised comparison of intravenous paracetamol and intravenous morphine for acute traumatic limb pain in the emergency department. *Emerg Med J* 2012; 29: pp. 37-39
[View In Article](#) | [Cross Ref](#)
179. Craig M., Jeavons R., Probert J., et al: Randomised comparison of intravenous paracetamol and intravenous morphine for acute traumatic limb pain in the emergency department. *Emerg Med J* 2012; 29: pp. 37-39
[View In Article](#) | [Cross Ref](#)
180. Cromhout A.: Ketamine: Its use in the emergency department. *Emerg Med* 2003; 15: pp. 155-159
[View In Article](#) | [Cross Ref](#)

181. Curran Jr WJC, Scott CB, Weinstein AS, Martin LA, Nelson JS, Phillips TL, Murray K, Fischbach AJ, Yakar D, Schwade JG (1993) Survival comparison of radiosurgery-eligible and -ineligible malignant glioma patients treated with hyperfractionated radiation therapy and carmustine: a report of Radiation Therapy Oncology Group 83-02. *J Clin Oncol* 11(5):857–862. <https://doi.org/10.1200/jco.1993.11.5.857> CrossRef Google Scholar
182. Curtis KM, Henriques HF, Fanciullo G, Reynolds CM, Suber F. A fentanyl-based pain management protocol provides early analgesia for adult trauma patients. *J Trauma*. 2007;63:819–826. [PubMed]
183. Dale O., Hjortkjaer R., and Kharasch E.D.: Nasal administration of opioids for pain management in adults. *Acta Anaesthesiol Scand* 2002; 46: pp. 759-770
[View In Article](#) | [Cross Ref](#)
184. D'Arcy Y.: Targeted topical analgesics for acute pain. *Pain Med News* 2014; 12: pp. 56-63
[View In Article](#)
185. Dart R.C., and Green J.L.: The prescription paradox of acetaminophen safety. *Pharmacoepidemiol Drug Saf* 2016 May; 25: pp. 599-601
[View In Article](#) | [Cross Ref](#)
186. Dart R.C., Surratt H.L., Cicero T.J., et al: Trends in opioid analgesic abuse and mortality in the United States. *N Engl J Med* 2015; 372: pp. 241-248
[View In Article](#) | [Cross Ref](#)
187. Davenport K., and Waine E.: The role of non-steroidal anti-inflammatory drugs in renal colic. *Pharmaceuticals* 2010; 3: pp. 1304-1310
[View In Article](#) | [Cross Ref](#)
188. Dela Cruz Ubaldo C, Hall NS, Le B. Postmarketing review of intravenous acetaminophen dosing based on Food and Drug Administration prescribing guidelines. *Pharmacotherapy* 2014;34:34S-39S.
189. Derry S., Conaghan P., Da Silva J.A., Wiffen P.J., and Moore R.A.: Topical NSAIDs for chronic musculoskeletal pain in adults. *Cochrane Database Syst Rev* 2016; undefined: pp. CD007400
[View In Article](#)
190. Derry S., Moore R.A., Gaskell H., McIntyre M., and Wiffen P.J.: Topical NSAIDs for acute musculoskeletal pain in adults. *Cochrane Database Syst Rev* 2015; undefined: pp. CD007402
[View In Article](#)
191. Dewhirst E., Frazier W.J., Leder M., Fraser D.D., and Tobias J.D.: Cardiac arrest following ketamine administration for rapid sequence intubation. *J Int Care Med* 2013; 28: pp. 375-379
[View In Article](#) | [Cross Ref](#)
192. Di Iorio D, Henley E, Doughty A. A survey of primary care physician practice patterns and adherence to acute low back problem guidelines. *Arch Fam Med*. 2000; 9:1015–21.
193. Di Iorio D, Henley E, Doughty A. A survey of primary care physician practice patterns and adherence to acute low back problem guidelines. *Arch Fam Med*. 2000;9:1015–1021. [PubMed]

194. Dickenson A.H.: NMDA receptor antagonists: interactions with opioids. *Acta Anaesthesiol Scand* 1997; 41: pp. 112-115
[View In Article](#) | [Cross Ref](#)
195. Dillard J.N., and Knapp S.: Complementary and alternative pain therapy in the emergency department. *Emerg Med Clin North Am* 2005; 23: pp. 529-549
[View In Article](#) | [Cross Ref](#)
196. Dordoni P., Ventura M.D., Stefanelli A., et al: Effect of ketorolac, ketoprofen and nefopam on platelet function. *Anaesthesia* 1994; 49: pp. 1046-1049
[View In Article](#) | [Cross Ref](#)
197. Doverty M., White J.M., Somogyi A.A., Bochner F., Ali R., and Ling W.: Hyperalgesic responses in methadone maintenance patients. *Pain* 2001 Feb 1; 90: pp. 91-96
[View In Article](#) | [Cross Ref](#)
198. Dowell D., Haegerich T.M., and Chou R.: CDC guideline for prescribing opioids for chronic pain—United States, 2016. *MMWR Recomm Rep* 2016; 65: pp. 1-49
[View In Article](#)
199. Dowell D., Haegerich T.M., and Chou R.: CDC guideline for prescribing opioids for chronic pain - United States, 2016. *MMWR Recomm Rep* 2016 Mar 18; 65: pp. 1-49
[View In Article](#)
200. Downing S.M., and Yudkowsky R.: Introduction to assessment in the health professions. In Downing S.M., and Yudkowsky R. (eds): *Assessment in health professions education*, 1st edn. New York: Routledge, 2009. pp. 1-20
[View In Article](#)
201. Dreher C, Habermehl D, Jakel O, Combs SE (2017) Effective radiotherapeutic treatment intensification in patients with pancreatic cancer: higher doses alone, higher RBE or both? *Radiat Oncol* 12(1):203. <https://doi.org/10.1186/s13014-017-0945-2>
[CrossRef](#)
[PubMed](#)
[PubMed Central](#)
[Google Scholar](#)
202. Drendel AL, Brousseau DC, Gorelick MH. Pain assessment for pediatric patients in the emergency department. *Pediatrics*. 2006;117:1508–1511. [\[PubMed\]](#)
203. Ducharme J. Emergency pain management: a Canadian Association of Emergency Physicians (CAEP) consensus document. *J Emerg Med* 1994; 12:855–66.
204. Ducharme J. Non-opioid pain medications to consider for emergency department patients. Available at: <http://www.acepnow.com/article/non-opioid-pain-medications-consider-emergency-department-patients/>. Accessed February 11, 2015.
[View In Article](#)
205. Duffy B.: Low-dose Ketamine Infusion Appears Safe Even Without Continuous Hemodynamic Monitoring. <http://www.anesthesiologynews.com/Clinical-Anesthesiology/Article/05-16/Low-Dose-Ketamine-Infusion-Appears-Safe-Even-Without-Continuous-Hemodynamic-Monitoring/36136>
[View In Article](#)
206. Dunn K.M., Saunders K.W., Rutter C.M., et al: Opioid prescriptions for chronic pain and overdose: a cohort study. *Ann Intern Med* Jan 2010; 152:
[View In Article](#)

207. Dunn L.K., and Durieux M.E.: Perioperative use of intravenous lidocaine. *Anesthesiology* 2017; 126: pp. 729-737
[View In Article](#) | [Cross Ref](#)
208. DuPen A, Shen D, Ersek M. Mechanisms of opioid-induced tolerance and hyperalgesia. *Pain Manag Nurs* 2007;8:113-21.
209. Durante M, Loeffler JS (2010) Charged particles in radiation oncology. *Nat Rev Clin Oncol* 7(1):37-43. <https://doi.org/10.1038/nrclinonc.2009.183>
[CrossRef](#)[PubMed](#)[Google Scholar](#)
210. Duthie D.J., and Nimmo W.S.: Adverse effects of opioid analgesic drugs. *Br J Anaesth* 1987; 59: pp. 61-77
[View In Article](#) | [Cross Ref](#)
211. Ead 2003.
[View In Article](#)
212. Ead H.: Review of laryngospasm and noncardiogenic pulmonary edema. *Dynamics* 2003; 14: pp. 9-12
[View In Article](#)
213. Edalat L, Stegen B, Klumpp L, Haehl E, Schilbach K, Lukowski R, Kuhnle M, Bernhardt G, Buschauer A, Zips D, Ruth P, Huber SM (2016) BK K + channel blockade inhibits radiation-induced migration/brain infiltration of glioblastoma cells. *Oncotarget* 7(12):14259–14278. <https://doi.org/10.18632/oncotarget.7423>
[CrossRef](#)[PubMed](#)[PubMed Central](#)[Google Scholar](#)
214. Eder SC, Sloan EP, Todd K. Documentation of ED patient pain by nurses and physicians. *Am J Emerg Med.* 2003;21:253–257. [\[PubMed\]](#)
215. Efron B., and Tibshirani R.: An introduction to the bootstrap. Boca Raton, FL: Chapman & Hall/CRC, 1998.
[View In Article](#)
216. Eide Stubhaug A., Oye I., and Breivik H.: Continuous subcutaneous administration of the . *Pain* 1995; 61: pp. 221-228
[View In Article](#) | [Cross Ref](#)
217. Eide Stubhaug A., Oye I., and Breivik H.: Continuous subcutaneous administration of the . *Pain* 1995; 61: pp. 221-228
[View In Article](#) | [Cross Ref](#)
218. Eisenhart A.W., Gaeta T.J., and Yens D.P.: Osteopathic manipulative treatment in the emergency department for patients with acute ankle injuries. *J Am Osteopath Assoc* 2003; 103: pp. 417-421
[View In Article](#)
219. Eken C, Serinken M, Elicabuk H, Uyanik E, Erdal M. Intravenous paracetamol versus dexketoprofen versus morphine in acute mechanical low back pain in the emergency department: a randomized double-blind controlled trial. *Emerg Med J* 2014;31:177-81.
220. Elenbaas RM, Waeckerle JF, McNabney WK. The clinical pharmacist in emergency medicine. *Am J Hosp Pharm.* 1977; 34:843-6.
221. Elenbaas RM. Role of the pharmacist in providing clinical pharmacy services in the emergency department. *Can J Hosp Pharm.* 1978; 31:123-5.
222. Ely E.W., Truman B., Shintani A., et al: Monitoring sedation status over time in ICU patients: reliability and validity of the Richmond Agitation-Sedation Scale (RASS).

JAMA 2003; 289: pp. 2983-2991

[View In Article](#) | [Cross Ref](#)

223. Ely E.W., Truman B., Shintani A., Thomason J.W., Wheeler A.P., Gordon S., et al: Monitoring sedation status over time in ICU patients: reliability and validity of the Richmond Agitation-Sedation Scale (RASS). JAMA 2003; 289: pp. 2983-2991
[View In Article](#) | [Cross Ref](#)
224. Emergency Nurses Association. RN procedural sedation rules. www.ena.org/government/State/Documents/RN_ProceduralSedationRules.pdf (accessed 2015 May 13).
225. Epstein R.M., and Hundert E.M.: Defining and assessing professional competence. JAMA 2002; 287: pp. 226-235
[View In Article](#) | [Cross Ref](#)
226. European Society for Emergency Medicine website. European curriculum for emergency medicine. Available at: <http://www.eusem.org/curriculumofem>. Accessed November 28, 2015.
[View In Article](#)
227. Evans E., Turley N., Robinson N., et al: Randomised controlled trial of patient controlled analgesia compared with nurse delivered analgesia in an emergency department. Emerg Med J 2005; 22: pp. 25-29
[View In Article](#) | [Cross Ref](#)
228. Fairbanks RJ, Hays DP, Webster DF et al. Clinical pharmacy services in an emergency department. Am J Health-Syst Pharm. 2004; 61:934-7.
229. Fairbanks RJ, Hildebrand JM, Kolstee KE et al. Medical and nursing staff highly value clinical pharmacists in the emergency department. Emerg Med J. 2007; 24:716-8.
230. Farnia M., et al: Comparison of intranasal ketamine versus IV morphine in reducing pain in patients with renal colic. Am J Emerg Med 2017; 35: pp. 434-437
[View In Article](#) | [Cross Ref](#)
231. Farnia MR, Babaei R, Shirani F, Momeni M, Hajimaghsoodi M, Vahidi E, et al. Analgesic effect of paracetamol combined with low-dose morphine versus morphine alone on patients with biliary colic:a double blind, randomized controlled trial. World J Emerg Med. 2016;7(1):25-9. [PMC free article] [PubMed]
232. Fathi M., Zare M.A., Bahmani H.R., and Zehtabchi S.: Comparison of oral oxycodone and naproxen in soft tissue injury pain control: a double-blind randomized clinical trial. Am J Emerg Med 2015 Sep; 33: pp. 1205-1208
[View In Article](#) | [Cross Ref](#)
233. Fein D.M., Avner J.R., Scharbach K., Manwani D., and Khine H.: Intranasal fentanyl for initial treatment of vaso-occlusive crisis in sickle cell disease. Pediatr Blood Cancer 2017; 64:
[View In Article](#)
234. Fenton J.J., Jerant A.F., Bertakis K.D., and Franks P.: The cost of satisfaction: a national study of patient satisfaction, health care utilization, expenditures, and mortality. Arch Intern Med 2012; 172: pp. 405-411
[View In Article](#) | [Cross Ref](#)
235. Finkel J., Pestieau S., and Quezado Z.: Ketamine as an adjuvant for treatment of cancer pain in children and adolescents. J Pain 2007; 8: pp. 515-521
[View In Article](#) | [Cross Ref](#)

236. Firouzian A., Alipour A., Rashidian Dezfooli H., et al: Does lidocaine as an adjuvant to morphine improve pain relief in patients presenting to the ED with acute renal colic? A double-blind, randomized controlled trial. *Am J Emerg Med* 2016 Mar; 34: pp. 443-448
[View In Article](#) | [Cross Ref](#)
237. Firouzian A., Alipour A., Rashidian Dezfooli H., et al: Does lidocaine as an adjuvant to morphine improve pain relief in patients presenting to the ED with acute renal colic? a double-blind, randomized controlled trial. *Am J Emerg Med* 2016; 34: pp. 443-448
[View In Article](#) | [Cross Ref](#)
238. Fischer B., Jones W., Urbanoski K., et al: Correlations between prescription opioid analgesic dispensing levels and related mortality and morbidity in Ontario, Canada, 2005-2011. *Drug Alcohol Rev* 2014; 33: pp. 19-26
[View In Article](#) | [Cross Ref](#)
239. Fischer B., Jones W., Urbanoski K., Skinner R., and Rehm J.: Correlations between prescription opioid analgesic dispensing levels and related mortality and morbidity in Ontario, Canada, 2005–2011. *Drug Alcohol Rev* 2014 Jan; 33: pp. 19-26
[View In Article](#) | [Cross Ref](#)
240. Fitch R.W., and Kuhn J.E.: Intraarticular lidocaine versus intravenous procedural sedation with narcotics and benzodiazepines for reduction of the dislocated shoulder: a systematic review. *Acad Emerg Med* 2008; 15: pp. 703-708
[View In Article](#) | [Cross Ref](#)
241. Fitzek MM, Thornton AF, Harsh G, Rabinov JD, Munzenrider JE, Lev M, Ancukiewicz M, Bussiere M, Hedley-Whyte ET, Hochberg FH, Pardo FS (2001) Dose-escalation with proton/photon irradiation for Daumas-Duport lower-grade glioma: results of an institutional phase I/II trial. *Int J Radiat Oncol Biol Phys* 51(1):131–137[CrossRef](#)[PubMed](#)[Google Scholar](#)
242. Fitzpatrick B.M., and Mullins M.E.: Intravenous lidocaine for the treatment of acute pain in the emergency department. *Clin Exp Emerg Med* 2016; 3: pp. 105-108
[View In Article](#) | [Cross Ref](#)
243. Fletcher D., and Martinez V.: Opioid-induced hyperalgesia in patients after surgery: a systematic review and a meta-analysis. *Br J Anaesth* 2014 Jun; 112: pp. 991-1004
[View In Article](#) | [Cross Ref](#)
244. Folts 1975.
[View In Article](#)
245. Folts J.D., Alfonso S., and Rowe G.G.: Systemic and coronary hemodynamic effects of ketamine in intact anaesthetized and unanaesthetized dogs. *Br J Anaesth* 1975; 47: pp. 686-694
[View In Article](#) | [Cross Ref](#)
246. Food and Drug Administration. Toradol® oral (ketorolac tromethamine tablets). Roche Laboratories Inc. Available at:
http://www.accessdata.fda.gov/drugsatfda_docs/label/2013/019645s019lbl.pdf. Accessed February 17, 2016.
[View In Article](#)

247. Ford L.E.: Acute hypertensive pulmonary edema: a new paradigm. *Can J Physiol Pharmacol* 2010; 88: pp. 9-13
[View In Article](#) | [Cross Ref](#)
248. Fosnocht DE, Heaps ND, Swanson ER. Patient expectations for pain relief in the ED. *Am J Emerg Med.* 2004;22:286–288. [\[PubMed\]](#)
249. Fosnocht DE, Swanson ER, Barton ED. Changing attitudes about pain and pain control in emergency medicine. *Emerg Med Clin North Am.* 2005;23:297–306. [\[PubMed\]](#)
250. Fosnocht DE, Swanson ER. Use of a triage pain protocol in the ED. *Am J Emerg Med.* 2007;25:791–793. [\[PubMed\]](#)
251. Frei SP, Bond WF, Bazuro RK, Richardson DM, Sierzega GM, Wasser TE. Is early analgesia associated with delayed treatment of appendicitis? *Am J Emerg Med.* 2008;26:176–180. [\[PubMed\]](#)
252. Friedman B.W., Adewunmi V., Campbell C., et al: A randomized trial of IV ketorolac versus IV metoclopramide plus diphenhydramine for tension-type and all non-migraine, non-cluster recurrent headaches. *Ann Emerg Med* 2013; 62: pp. 311-318
[View In Article](#) | [Cross Ref](#)
253. Friedman B.W., Dym A.A., Davitt M., Holden L., Solorzano C., Esses D., et al: Naproxen with cyclobenzaprine, oxycodone/acetaminophen, or placebo for treating acute low back pain: a randomized clinical trial. *JAMA* 2015 Oct 20; 314: pp. 1572-1580
[View In Article](#) | [Cross Ref](#)
254. Fuentes EF, Kohn MA, Neighbor ML. Lack of association between patient ethnicity or race and fracture analgesia. *Acad Emerg Med.* 2002;9:910–915. [\[PubMed\]](#)
255. Furyk J.S., Grabowski W.J., and Black L.H.: Nebulized fentanyl versus intravenous morphine in children with suspected limb fractures in the emergency department: a randomized controlled trial. *Emerg Med Australas* 2009; 21: pp. 203-209
[View In Article](#) | [Cross Ref](#)
256. Furyk J.S., Grabowski W.J., and Black L.H.: Nebulized fentanyl versus intravenous morphine in children with suspected limb fractures in the emergency department: a randomized controlled trial. *Emerg Med Australas* 2009; 21: pp. 203-209
[View In Article](#) | [Cross Ref](#)
257. Galinski M, Dolveck F, Combes X, et al. Management of severe acute pain in emergency settings: ketamine reduces morphine consumption. *Am J Emerg Med* 2007;25:385–90.
258. Galinski M., Dolveck F., Combes X., et al: Management of severe acute pain in emergency settings: ketamine reduces morphine consumption. *Am J Emerg Med* 2007; 25: pp. 385-390
[View In Article](#) | [Cross Ref](#)
259. Galinski M., Dolveck F., Combes X., et al: Management of severe acute pain in emergency settings: ketamine reduces morphine consumption. *Am J Emerg Med* 2007; 25: pp. 385-390
[View In Article](#) | [Cross Ref](#)
260. Galinski M., Dolveck F., Combes X., et al: Management of severe acute pain in emergency settings: ketamine reduces morphine consumption. *Am J Emerg Med* 2007; 25: pp. 385-390
[View In Article](#) | [Cross Ref](#)

261. Galinski M., Dolveck F., Combes X., Limoges V., Smail N., Pommier V., et al: Management of severe acute pain in emergency settings: ketamine reduces morphine consumption. *Am J Emerg Med* 2007;May; 25: pp. 385-390
[View In Article](#) | [Cross Ref](#)
262. Galinski M., Dolveck F., Combes X., Limoges V., Smail N., Pommier V., et al: Management of severe acute pain in emergency settings: ketamine reduces morphine consumption. *Am J Emerg Med* 2007; 25: pp. 385-390
[View In Article](#) | [Cross Ref](#)
263. Galinski M.M., Dolveck F.M., Combes X.M., et al: Management of severe acute pain in emergency settings: ketamine reduces morphine consumption. *Am J Emerg Med* 2007; 25: pp. 385-390
[View In Article](#) | [Cross Ref](#)
264. Gallagher EJ, Esses D, Lee C, Lahn M, Bijur PE. Randomized clinical trial of morphine in acute abdominal pain. *Ann Emerg Med*. 2006;48:150–160. [[PubMed](#)]
265. Gallagher J.E., Blauth J., and Fornadley J.A.: Perioperative ketorolac tromethamine and postoperative hemorrhage in cases of tonsillectomy and adenoidectomy. *Laryngoscope* 1995; 105: pp. 606-609
[View In Article](#) | [Cross Ref](#)
266. Gallagher JE, Blauth J, Fornadley JA. Perioperative Ketorolac tromethamine and postoperative hemorrhage in cases of tonsillectomy and adenoidectomy. *Laryngoscope*. 1995;105:606–609. [[PubMed](#)]
267. Gamis A.S., Knapp J.F., and Glenski J.A.: Nitrous oxide analgesia in a pediatric emergency department. *Ann Emerg Med* 1989; 18: pp. 177-181
[View In Article](#) | [Cross Ref](#)
268. Gandhi S.K., Powers J.C., Normeir A.M., Fowle K., Kitzman D.W., Rankin K.M., et al: The pathogenesis of acute pulmonary edema associated with hypertension. *N Engl J Med* 2001; 344: pp. 17-22
[View In Article](#) | [Cross Ref](#)
269. Gao M, Rejaei D, Liu H. Ketamine use in current clinical practice. *Acta Pharmacol Sin*. 2016;37(7):865–72. Epub 2016 Mar 28. [[PMC free article](#)] [[PubMed](#)]
270. Gao M, Rejaei D, Liu H. Ketamine use in current clinical practice. *Acta Pharmacol Sin*. 2016;37(7):865–72. [[PMC free article](#)] [[PubMed](#)]
271. Gao M., Rejaei D., and Liu H.: Ketamine use in current clinical practice. *Acta Pharmacol Sin* 2016; 37: pp. 865-872
[View In Article](#) | [Cross Ref](#)
272. Gao M., Rejaei D., and Liu H.: Ketamine use in current clinical practice. *Acta Pharmacol Sin* 2016; 37: pp. 865-872
[View In Article](#) | [Cross Ref](#)
273. García Rodríguez L.A., Cattaruzzi C., Troncon M.G., et al: Risk of hospitalization for upper gastrointestinal tract bleeding associated with ketorolac, other nonsteroidal anti-inflammatory drugs, calcium antagonists, and other antihypertensive drugs. *Arch Intern Med* 1998; 158: pp. 33-39
[View In Article](#) | [Cross Ref](#)
274. García Rodríguez LA, Cattaruzzi C, Troncon MG, Agostinis L. Risk of hospitalization for upper gastrointestinal tract bleeding associated with ketorolac, other nonsteroidal anti-inflammatory drugs, calcium antagonists, and other antihypertensive drugs. *Arch Intern Med*. 1998;158(1):33–9. [[PubMed](#)]

275. Gaskin DJ, Richard P. The economic costs of pain in the United States. In: relieving pain in America: a blueprint for transforming prevention, care, education, and research. 2011. Available at: http://www.ncbi.nlm.nih.gov/books/NBK91497/pdf/Bookshelf_NBK91497.pdf. Accessed February 6, 2016.
[View In Article](#)
276. Gauntlet-Gilbert J., Rodham K., Jordan A., et al: Emergency department staff attitudes toward people presenting in chronic pain: a qualitative study. *Pain Med* 2015; 16: pp. 2065-2074
[View In Article](#) | [Cross Ref](#)
277. Gibson S.J., and Farrell M.: What is different about pain in older people? *Rev Analgesia* 2004; 8: pp. 23-37
[View In Article](#) | [Cross Ref](#)
278. Gibson S.J.: Older Persons' Pain: what can we learn? *Pain Clin Updat* 2006; 14: pp. 1-4
[View In Article](#)
279. Gibson S.J.: Pain and ageing: a comparison of the pain experience over the adult life span. *Prog Pain Res Manage* 2003; 24: pp. 767-790
[View In Article](#)
280. Godwin S.A., Burton J.H., Gerardo C.J., Hatten B.W., Mace S.E., Silvers S.M., et al: Clinical Policy: Procedural sedation and analgesia in the Emergency Department. *Ann Emerg Med* 2014; 63: pp. 247-258
[View In Article](#) | [Cross Ref](#)
281. Goldman RD, Narula N, Klein-Kremer, Finkelstein Y, Rogovik AL. Predictors for opioid analgesia administration in children with abdominal pain presenting to the emergency department. *Clin J Pain*. 2008;24:11–15. [[PubMed](#)]
282. Goldman RD, Narula N, Klein-Kremer, Finkelstein Y, Rogovik AL. Predictors for opioid analgesia administration in children with abdominal pain presenting to the emergency department. *Clin J Pain*. 2008; 24:11-5.
283. Goltser A, Soleyman-Zomalan E, Kresch F, et al. Short (low-dose) ketamine infusion for managing acute pain in the ED: case-report series. *Am J Emerg Med*. 2015;33(4):601.e5-7. [[PubMed](#)]
284. Goltser A, Soleyman-Zomalan E, Kresch F, Motov S. Short (low-dose) ketamine infusion for managing acute pain in the ED:case-report series. *Am J Emerg Med*. 2015;33(4):601.e5-7. [[PubMed](#)]
285. Goltser A., Soleyman-Zomalan E., Kresch F., and Motov F.: Short (low-dose) ketamine infusion for managing acute pain in the ED: case-report series. *Am J Emerg Med* 2015; 33: pp. 601.e5-601.e7
[View In Article](#) | [Cross Ref](#)
286. Goltser A., Soleyman-Zomalan E., Kresch F., and Motov S.: Short (low-dose) ketamine infusion for managing acute pain in the ED: case-report series. *Am J Emerg Med* 2015; 33: pp. 601.e5-7
[View In Article](#)
287. Goltser A., Soleyman-Zomalan E., Kresch F., and Motov S.: Short (low-dose) ketamine infusion for managing acute pain in the ED: case-report series. *Am J Emerg*

Med 2015 Apr; 33:

[View In Article](#)

288. Goltser A., Soleyman-Zomalan E., Kresch F., et al: Short (low-dose) ketamine infusion for managing acute pain in the ED: case-report series. Am J Emerg Med 2015; 33:

[View In Article](#)

289. Goltser A., Soleyman-Zomalan E., Kresch F., et al: Short (low-dose) ketamine infusion for managing acute pain in the ED: case-report series. Am J Emerg Med 2015; 33: pp. 601.e5-601.e7

[View In Article](#) | [Cross Ref](#)

290. Golzari S.E., Soleimanpour H., Mahmoodpoor A., et al: Lidocaine and pain management in the emergency department: a review article. Anesth Pain Med 2014; 4: pp. e15444

[View In Article](#)

291. Golzari S.E., Soleimanpour H., Rahmani F., et al: Therapeutic approaches for renal colic in the emergency department: a review article. Anesth Pain Med 2014 Feb 13; 4:

[View In Article](#)

292. Golzari SE, Soleimanpour H, Mahmoodpoor A et al. Lidocaine and pain management in the emergency department: a review article. Anesth Pain Med. 2014; 4:e15444.

293. Goodacre SW, Roden RK. A protocol to improve analgesia use in the accident and emergency department. J Accid Emerg Med. 1996;13:177–179. [[PMC free article](#)] [[PubMed](#)]

294. Gordh T., Karlsten R., and Kristensen J.: Intervention with spinal NMDA, adenosine and NO systems for pain modulation. Ann Med 1995; 27: pp. 229-234

[View In Article](#) | [Cross Ref](#)

295. Gorlin A.W., Rosenfeld D.M., and Ramakrishna H.: Intravenous sub-anesthetic ketamine for perioperative analgesia. J Anaesthesiol Clin Pharmacol 2016; 32: pp. 160-167

[View In Article](#) | [Cross Ref](#)

296. Grading quality of evidence and strength of recommendations. BMJ 2004; 328:

[View In Article](#)

297. Graudins A., Meek R., Egerton-Warburton D., et al: The PICHFORK (Pain in Children Fentanyl or Ketamine) trial: a randomized controlled trial comparing intranasal ketamine and fentanyl for the relief of moderate to severe pain in children with limb injuries. Ann Emerg Med 2015; 65: pp. 248-254

[View In Article](#) | [Cross Ref](#)

298. Green 1990.

[View In Article](#)

299. Green 1990.

[View In Article](#)

300. Green 2000.

[View In Article](#)

301. Green 2000.

[View In Article](#)

302. Green 2000.
[View In Article](#)
303. Green 2000.
[View In Article](#)
304. Green 2000.
[View In Article](#)
305. Green R, Bulloch B, Kabani A, Hnacock BJ, Tenenbein M. Early analgesia for children with acute abdominal pain. *Pediatrics*. 2005;116:978–983. [PubMed]
306. Green S.M., and Johnson N.E.: Ketamine sedation for pediatric procedures: Part 2, Review and implications. *Ann Emerg Med* 1990; 19: pp. 1033-1046
[View In Article](#) | [Cross Ref](#)
307. Green S.M., and Johnson N.E.: The Semantics of Ketamine. *Ann Emerg Med* 2000; 36: pp. 480-482
[View In Article](#)
308. Green S.M., and Li J.: Ketamine in Adults What Emergency Physicians Need to Know about Patient Selection and Emergence Reactions. *Acad Emerg Med* 2000; 7: pp. 278-281
[View In Article](#) | [Cross Ref](#)
309. Green S.M., Roback M.G., Krauss B., Brown L., McGlone R.G., Agrawal D., et al: Predictors of airway and respiratory adverse events with ketamine sedation in the emergency department: an individual-patient data meta-analysis of 8,282 children. *Ann Emerg Med* 2009; 54: pp. 158-168
[View In Article](#) | [Cross Ref](#)
310. Green S.M.: There is oligo-evidence for oligoanalgesia. *Ann Emerg Med* 2012; 60: pp. 212-214
[View In Article](#) | [Cross Ref](#)
311. Green SM, Li J. Ketamine in adults: what emergency physicians need to know about patient selection and emergence reactions. *Acad Emerg Med* 2000;7:278–81.
312. Green SM, Roback MG, Kennedy RM, Krauss B. Clinical practice guideline for emergency department ketamine dissociative sedation: 2011 update. *Ann Emerg Med* 2011;57:449–61.
313. Greenwood-Ericksen M.B., Poon S.J., Nelson L.S., Weiner S.G., and Schuur J.D.: Best practices for prescription drug monitoring programs in the emergency department setting: results of an expert panel. *Ann Emerg Med* 2016; 67: pp. 755-764
[View In Article](#) | [Cross Ref](#)
314. Greer I.A.: Effects of ketorolac tromethamine on hemostasis. *Pharmacotherapy* 1990; 10: pp. 71S-76S
[View In Article](#)
315. Grissa MH, Claessens YE, Bouida W, et al. Paracetamol vs piroxicam to relieve pain in renal colic. Results of a randomized controlled trial. *Am J Emerg Med* 2011;29:203–6.
316. Guirimand F.M., Dupont X.M., Brasseur L.M., et al: The effects of ketamine on the temporal summation (wind-up) of the RIII nociceptive flexion reflex and pain in humans. *Anesth Analg* 2000; 90: pp. 408-414
[View In Article](#)

317. Gurnani A, Sharma PK, Rautela RS, Bhattacharya A. Analgesia for acute musculoskeletal trauma: low-dose subcutaneous infusion of ketamine. *Anaesth Intensive Care* 1996;24:32-6.
318. Gurnani A., Sharma P.K., Rautela R.S., and Bhattacharya A.: Analgesia for acute musculoskeletal trauma: low-dose subcutaneous infusion of ketamine. *Anaesth Intensive Care* 1996; 24: pp. 32-36
[View In Article](#)
319. Guru V, Dubinsky I. The patient vs caregiver perception of acute pain in the emergency department. *J Emerg Med*. 2000;18:7–12. [PubMed]
320. Haas DA, Harper DG. Ketamine: a review of its pharmacologic properties and use in ambulatory anesthesia. *Anesth Prog* 1992;39:61-8.
321. Haines L., Dickman E., Ayvazyan S., et al: Ultrasound-guided fascia iliaca compartment block for hip fractures in the emergency department. *J Emerg Med* 2012; 43: pp. 692-697
[View In Article](#) | [Cross Ref](#)
322. Hartrick CT, Kovan JP, Shapiro S. The numeric rating scale for clinical pain measurement: a ratio measure? *Pain Pract*. 2003; 3:310-6.
323. Hazhir S., Badr Y.A., and Darabi J.N.: Comparison of intranasal desmopressin and intramuscular tramadol versus pethidine in patients with renal colic. *Urol J* 2010; 7: pp. 148-151
[View In Article](#)
324. Hedegaard M.D., Chen M.S., and Warner M.: Drug-poisoning deaths involving heroin: United States, 2000–2013. *Natl Cent Health Stat Data Brief* 2015; 190: pp. 1-8
[View In Article](#)
325. Hernández-Díaz S., and García-Rodríguez L.A.: Epidemiologic assessment of the safety of conventional nonsteroidal anti-inflammatory drugs. *Am J Med* 2001; 110: pp. 20S-27S
[View In Article](#)
326. Herr K., and Titler M.: Acute pain assessment and pharmacological management practices for the older adult with a hip fracture: review of ED trends. *J Emerg Nurs* 2009 Jul; 35: pp. 312-320
[View In Article](#) | [Cross Ref](#)
327. Herring AA, Ahern T, Stone MB, Frazee BW. Emerging applications of low-dose ketamine for pain management in the ED. *Am J Emerg Med* 2013;31:416-9.
328. Higgins J.P., Altman D.G., Gotzsche P.C., et al: The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011; 343: pp. d5928
[View In Article](#) | [Cross Ref](#)
329. Higgins J.P., and Green S.: Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. Denmark: Cochrane Collaboration, 2011.
[View In Article](#)
330. Hill G.E., Anderson J.L., and Whitten C.W.: Ketamine inhibits agonist-induced cAMP accumulation increase in human airway smooth muscle cells. *Can J Anesth* 1999; 46: pp. 1172-1177
[View In Article](#) | [Cross Ref](#)

331. Himmelseher S, Durieux ME. Ketamine for perioperative pain management. *Anesthesiology* 2005;102:211-20.
332. Hina N., Fletcher D., Poindessous-Jazat F., and Martinez V.: Hyperalgesia induced by low-dose opioid treatment before orthopaedic surgery: an observational case-control study. *Eur J Anaesthesiol* 2015 Apr; 32: pp. 255-261
[View In Article](#) | [Cross Ref](#)
333. Hirota Y, Masunaga S, Kondo N, Kawabata S, Hirakawa H, Yajima H, Fujimori A, Ono K, Kuroiwa T, Miyatake S (2014) High linear-energy-transfer radiation can overcome radioresistance of glioma stem-like cells to low linear-energy-transfer radiation. *J Radiat Res* 55(1):75–83. <https://doi.org/10.1093/jrr/rrt095> [CrossRef](#)[PubMed](#)[Google Scholar](#)
334. Hocking G, Cousins MJ. Ketamine in chronic pain management: an evidence-based review. *Anesth Analg* 2003;97:1730–9.
335. Hoffman B.M., Papas R.K., Chatkoff D.K., and Kerns R.D.: Meta-analysis of psychological interventions for chronic low back pain. *Health Psychol* 2007; 26: pp. 1-9
[View In Article](#) | [Cross Ref](#)
336. Holdgate A., and Pollock T.: Systematic review of the relative efficacy of non-steroidal anti-inflammatory drugs and opioids in the treatment of acute renal colic. *BMJ* 2004; 328: pp. 1401
[View In Article](#) | [Cross Ref](#)
337. Holdgate A., Asha S., Craig J., et al: Comparison of a verbal numeric rating scale with the visual analogue scale for the measurement of acute pain. *Emerg Med Australas* 2003; 15: pp. 441-446
[View In Article](#) | [Cross Ref](#)
338. Holdgate A., Asha S., Craig J., et al: Comparison of a verbal numeric rating scale with the visual analogue scale for the measurement of acute pain. *Emerg Med (Fremantle)* 2003; 15: pp. 441-446
[View In Article](#) | [Cross Ref](#)
339. Holdgate A., Asha S., Craig J., et al: Comparison of a verbal numeric rating scale with the visual analogue scale for the measurement of acute pain. *Emerg Med Australas* 2003; 15: pp. 441-446
[View In Article](#) | [Cross Ref](#)
340. Hollmann M.W., and Durieux M.E.: Local anesthetics and the inflammatory response: a new therapeutic indication? *Anesthesiology* 2000; 93: pp. 858-875
[View In Article](#) | [Cross Ref](#)
341. Hooten W.M., Lamer T.J., and Twyner C.: Opioid-induced hyperalgesia in community-dwelling adults with chronic pain. *Pain* 2015 Jun; 156: pp. 1145-1152
[View In Article](#)
342. Hooten W.M., St Sauver J.L., McGree M.E., Jacobson D.J., and Warner D.O.: Incidence and risk factors for progression from short-term to episodic or long-term opioid prescribing: a population-based study. *Mayo Clin Proc* 2015 Jul; 90: pp. 850-856
[View In Article](#) | [Cross Ref](#)
343. Hoppe J.A., Houghland J., Yaron M., et al: Prescription history of emergency department patients prescribed opioids. *West J Emerg Med* 2013; 14: pp. 247-252
[View In Article](#) | [Cross Ref](#)

344. Hoppe J.A., Kim H., and Heard K.: Association of emergency department opioid initiation with recurrent opioid use. Ann Emerg Med 2015 May; 65: pp. 493-499
[View In Article](#) | [Cross Ref](#)
345. Hoppe J.A., Kim H., and Heard K.: Association of emergency department opioid initiation with recurrent opioid use. Ann Emerg Med 2015; 65: pp. 493-499.e4
[View In Article](#) | [Cross Ref](#)
346. Hoppe J.A., Nelson L.S., Perrone J., and Weiner S.G.: Opioid prescribing in a cross section of US emergency departments. Ann Emerg Med 2015 Sep; 66: pp. 253-259
[View In Article](#) | [Cross Ref](#)
347. Hoppe J.A., Nelson L.S., Perrone J., and Weiner S.G.: Opioid prescribing in a cross section of US emergency departments. Ann Emerg Med 2015; 66: pp. 253-259.e1
[View In Article](#) | [Cross Ref](#)
348. Hoppe J.A., Nelson L.S., Perrone J., and Weiner S.G.: Prescribing Opioids Safely in the Emergency Department (POSED) Study Investigators. Opioid prescribing in a cross section of US emergency departments. Ann Emerg Med 2015; 66: pp. 253-259
[View In Article](#) | [Cross Ref](#)
349. Hoppe JA, Nelson LS, Weiner SG et al. Planning for a safer decade of ED analgesia. Presentation at 25th Society of Academic Emergency Medicine annual meeting. Dallas, TX; 2014 May 13–17.
350. Hosseininejad S.M., Amini Ahidashti H., Bozorgi F., et al: Efficacy and safety of combination therapy with ketorolac and morphine in patient with acute renal colic; a triple-blind randomized controlled clinical trial. Bull Emerg Trauma 2017 Jul; 5: pp. 165-170
[View In Article](#)
351. Hou LC, Veeravagu A, Hsu AR, Tse VC (2006) Recurrent glioblastoma multiforme: a review of natural history and management options. Neurosurg Focus 20(4):E5[CrossRef](#)[PubMed](#)[Google Scholar](#)
352. Hwang U, Richardson LD, Sonya TO, Morrison RS. The effect of emergency department crowding on the management of pain in older adults with hip fracture. J Am Geriatr Soc. 2006;54:270–275. [[PubMed](#)]
353. Idrees U, Clements E. The state of U.S. emergency care: a call to action for hospital pharmacists. Ann Pharmacother. 2006; 40:2251-3.
354. Institute of Medicine website. Relieving pain in America: a blueprint for transforming prevention, care, education, and research. Available at:
<http://www.iom.edu/Reports/2011/Relieving-Pain-in-America-A-Blueprint-for-Transforming-Prevention-Care-Education-Research.aspx>. Accessed November 28, 2015.
[View In Article](#)
355. Institute of Medicine. Relieving Pain in America: A Blueprint for Transforming Prevention, Care, Education, and Research. http://www.nap.edu/catalog.php?record_id=13172. Accessed April 8, 2016.
[View In Article](#)
356. International Association for the Study of Pain website. International Association for the Study of Pain has released a comprehensive pain curriculum. Available at: www.iasp-pain.org/Education/CurriculumDetail.aspx?ItemNumber=729. Accessed November 28, 2015.
[View In Article](#)

357. International Narcotics Control Board. https://www.incb.org/incb/en/narcotic-drugs/Technical_Reports/narcotic_drugs_reports.html Accessed July 20, 2016.
[View In Article](#)
358. Inturrisi C.E.: Clinical pharmacology of opioids for pain. Clin J Pain 2012; 18: pp. S3-13
[View In Article](#) | [Cross Ref](#)
359. Irifune M., Shimizu T., Nomoto M., and Fukuda T.: Ketamine-induced anesthesia involves the N-methyl-D-aspartate receptor-channel complex in mice. Brain Res 1992; 569: pp. 1-9
[View In Article](#) | [Cross Ref](#)
360. Jasani N.B., O'Conner R.E., and Bouzoukis J.K.: Comparison of hydromorphone and meperidine for ureteral colic. Acad Emerg Med 1994 Nov-Dec; 1: pp. 539-543
[View In Article](#) | [Cross Ref](#)
361. Javery K.B., Ussery T.W., Steger H.G., and Colclough G.W.: Comparison of morphine and morphine with ketamine for postoperative analgesia. Can J Anaesth 1996; 43: pp. 212
[View In Article](#) | [Cross Ref](#)
362. Jennings P.A., Cameron P., and Bernard S.: Ketamine as an analgesic in the pre-hospital setting: a systematic review. Acta Anaesthesiol Scand 2011; 55: pp. 638-643
[View In Article](#) | [Cross Ref](#)
363. Jennings P.A., Cameron P., and Bernard S.: Ketamine as an analgesic in the pre-hospital setting: a systematic review. Acta Anaesthesiol Scand 2011; 55: pp. 638-643
[View In Article](#) | [Cross Ref](#)
364. Jennings P.A., Cameron P., Bernard S., et al: Morphine and ketamine is superior to morphine alone for out-of-hospital trauma analgesia: a randomized controlled trial. Ann Emerg Med 2012; 59: pp. 497-503
[View In Article](#) | [Cross Ref](#)
365. Jennings P.A., Cameron P., Bernard S., Walker T., Jolley D., and Masci K.: Morphine and ketamine is superior to morphine alone for out-of-hospital trauma analgesia: a randomized controlled trial. Ann Emerg Med 2012; 59: pp. 6
[View In Article](#) | [Cross Ref](#)
366. Jennings P.P., Cameron P.M., Bernard S.M., et al: Morphine and ketamine is superior to morphine alone for out-of-hospital trauma analgesia: a randomized controlled trial. Ann Emerg Med 2012; 59: pp. 497-503
[View In Article](#) | [Cross Ref](#)
367. Jennings PA, Cameron P, Bernard S. Ketamine as an analgesic in the pre-hospital setting:a systematic review. Acta Anaesthesiol Scand. 2011;55(6):638–43. [PubMed]
368. Jennings PA, Cameron P, Bernard S. Ketamine as an analgesic in the pre-hospital setting: a systematic review. Acta Anaesthesiol Scand. 2011;55(6):638–43. [PubMed]
369. Johansson P., Kongstad P., and Johansson A.: The effect of combined treatment with morphine sulphate and low-dose ketamine in a prehospital setting. Scand J Trauma Resusc Emerg Med 2009; 17: pp. 61
[View In Article](#)
370. Johansson P., Kongstad P., and Johansson A.: The effect of combined treatment with morphine sulphate and low-dose ketamine in a prehospital

setting. Scand J Trauma Resusc Emerg Med 2009; 17: pp. 61

[View In Article](#)

371. Johansson P., Kongstad P., and Johansson A.: The effect of combined treatment with morphine sulphate and low-dose ketamine in a prehospital setting. Scand J Trauma Resusc Emerg Med 2009; 17: pp. 1-5

[View In Article](#)

372. Johnson J.C., and Atherton G.L.: Effectiveness of nitrous oxide in a rural EMS system. J Emerg Med 1991; 9: pp. 45-53

[View In Article](#)

373. Johnstone M.: The cardiovascular effects of ketamine in man. Anaesthesia 1976; 31: pp. 873-882

[View In Article](#) | [Cross Ref](#)

374. Joint Commission on the Accreditation of Healthcare Organizations. CAMH Revised Pain Management Standards. 2000. Cited on Jul 20, 2001. Available from <http://www.jcaho.org/standard/pmhap.html>.

375. Joint Commission. Sentinel event alert 2012.

www.jointcommission.org/assets/1/18/SEA_49_opioids_8@_12_final.pdf (accessed 2014 Nov 1).

376. Jones J.B.: Assessment of pain management skills in emergency medicine residents: the role of a pain education program. J Emerg Med 1999; 17: pp. 349-354

[View In Article](#) | [Cross Ref](#)

377. Jones JB. Assessment of pain management skills in emergency medicine residents: the role of a pain education program. *J Emerg Med*. 1999; 17:349-54.

378. Jones JB. Assessment of pain management skills in emergency medicine residents: the role of a pain education program. *J Emerg Med*. 1999;17:349-354. [PubMed]

379. Jones JS, Johnson K, McNinch M. Age as a risk factor for inadequate emergency department analgesia. *Am J Emerg Med*. 1996;14:157-160. [PubMed]

380. Jones S.F., and O'Donnell A.M.: Clinical pharmacology: traditional NSAIDs and selective COX-2 inhibitors. In Macintyre P.E., Walker S.M., and Rowbotham D.J. (eds): Clinical Pain Management (Acute Pain), 2nd ed. London: England: Hodder & Stoughton Limited, 2008. pp. 68-79
[View In Article](#)

381. Jones S.F., and O'Donnell A.M.: Clinical pharmacology: traditional NSAIDs and selective COX-2 inhibitors. In Macintyre P.E., Walker S.M., and Rowbotham D.J. (eds): Clinical Pain Management (Acute Pain), 2nd ed. London, England: Hodder & Stoughton Limited, 2008. pp. 79
[View In Article](#)

382. Jones S.F., and O'Donnell A.M.: Clinical Pharmacology: traditional NSAIDs and selective COX-2 inhibitors. In Macintyre P.E., Walker S.M., and Rowbotham D.J. (eds): Clinical pain management (acute pain), 2nd edition. London: Hodder & Stoughton Limited, 2008. pp. 68-79
[View In Article](#)

383. Jouguet-Lacoste J., La Colla L., Schilling D., et al: The use of intravenous infusion or single dose of low-dose ketamine for postoperative analgesia: a review of the current literature. *Pain Med* 2015; 16: pp. 383-403
[View In Article](#) | [Cross Ref](#)

384. Jung D., Mroszczak E.J., Wu A., et al: Pharmacokinetics of ketorolac and p-hydroxyketorolac following oral and intramuscular administration of ketorolac tromethamine. *Pharm Res* 1989; 6: pp. 62-65
[View In Article](#) | [Cross Ref](#)
385. Juurlink D.N., Dhalla I.A., and Nelson L.S.: Improving opioid prescribing: the New York City recommendations. *JAMA* 2013 Mar 6; 309: pp. 879-880
[View In Article](#) | [Cross Ref](#)
386. Juurlink D.N., Dhalla I.A., and Nelson L.S.: Improving opioid prescribing: The New York City recommendations. *JAMA* 2013; 309: pp. 879-880
[View In Article](#) | [Cross Ref](#)
387. Juurlink DN. Improving opioid prescribing: the New York City recommendations. *JAMA*. 2013; 309:879-80.
388. Kaboli PJ, Hoth AB, McClimon BJ et al. Clinical pharmacists and inpatient medical care. *Arch Intern Med*. 2006; 166:955-64.
389. Kain Z.N., Mayes L.C., Ferris C.A., Pakes J., and Schottenfeld R.: Cocaine-abusing parturients undergoing cesarean section: a cohort study. *Anesthesiology* 1996; 85: pp. 1028-1035
[View In Article](#) | [Cross Ref](#)
390. Kallet R.H., Daniel B.M., Gropper M., and Matthay M.A.: Acute pulmonary edema following upper airway obstruction: case reports and brief review. *Resp Care* 1998; 43: pp. 476-480
[View In Article](#)
391. Kapural L., Kapural M., Bensitel T., and Sessler D.I.: Opioid-sparing effect of intravenous outpatient ketamine infusions appears short-lived in chronic-pain patients with high opioid requirements. *Pain Physician* 2010; 13: pp. 389-394
[View In Article](#)
392. Kariman H., Majidi A., Amini A., et al: Nitrous oxide/oxygen compared with fentanyl in reducing pain among adults with isolated extremity trauma: a randomized trial. *Emerg Med Australas* 2011; 23: pp. 761-768
[View In Article](#) | [Cross Ref](#)
393. Kasuya A, Bauman J, Curtis RA et al. Clinical pharmacy on-call program in the emergency department. *Am J Emerg Med*. 1986; 4:464-7.
394. Kaye A.D., Banister R.E., Anwar M., Feng C.J., Kadowitz P.J., and Nossaman B.D.: Pulmonary vasodilation by ketamine is mediated in part by L-type calcium channels. *Anesth Analg* 1998 Oct; 87: pp. 956-962
[View In Article](#)
395. Kelly AM, Brumby C, Barnes C. Nurse-initiated, titrated intravenous opioid analgesia reduces time to analgesia for selected painful conditions. *CJEM*. 2005;7:149–154. [[PubMed](#)]
396. Kelly S., Johnson G.T., and Harbison R.D.: “Pressured to prescribe” The impact of economic and regulatory factors on South-Eastern ED physicians when managing the drug seeking patient. *J Emerg Trauma Shock* 2016; 9: pp. 58-63
[View In Article](#)
397. Kendall J.M., Reeves B.C., and Latter V.S.: Multicentre randomised controlled trial of nasal diamorphine for analgesia in children and teenagers with clinical features. *BMJ* 2001; 322: pp. 261-265
[View In Article](#) | [Cross Ref](#)

398. Kennedy RM, Porter FL, Miller JP, Jaffe DM. Comparison of fentanyl/midazolam with ketamine/midazolam for pediatric orthopedic emergencies. *Pediatrics* 1998;102:956–63.
399. Kil WJ, Tofilon PJ, Camphausen K (2012) Post-radiation increase in VEGF enhances glioma cell motility in vitro. *Radiat Oncol* 7:25. <https://doi.org/10.1186/1748-717X-7-25>
[CrossRef](#)
[PubMed](#)
[PubMed Central](#)
[Google Scholar](#)
400. Kim MK, Galustyan S, Sato TT, Bergholte J, Hennes HM. Analgesia for children with acute abdominal pain: a survey of pediatric emergency physicians and pediatric surgeons. *Pediatrics*. 2003;112:1122–1126. [\[PubMed\]](#)
401. Kim MK, Strait RT, Sato TT, Hennes HM. A randomized clinical trial of analgesia in children with acute abdominal pain. *Acad Emerg Med*. 2002;9:281–287. [\[PubMed\]](#)
402. King N.B., Fraser V., Boikos C., Richardson R., and Harper S.: Determinants of increased opioid-related mortality in the United States and Canada, 1990-2013: a systematic review. *Am J Public Health* 2014 Aug; 104: pp. e32-e42
[View In Article](#) | [Cross Ref](#)
403. Klaess CC, Jungquist CR. Current Ketamine Practice:Results of the 2016 American Society of Pain Management Nursing Survey on Ketamine. *Pain Manag Nurs*. 2018;19(3):222–9. [\[PubMed\]](#)
404. Kleihues P, Louis DN, Scheithauer BW, Rorke LB, Reifenberger G, Burger PC, Cavenee WK (2002) The WHO classification of tumors of the nervous system. *J Neuropathol Exp Neurol* 61(3):215–225 (**discussion 226–219**)
[CrossRef](#)
[PubMed](#)
[Google Scholar](#)
405. Kohn LT, Corrigan JM, Donaldson MS, eds. To err is human: building a safer health system. Washington, DC: National Academy Press; 1999.
406. Kohrs 1998.
[View In Article](#)
407. Kohrs R., and Durieux M.E.: Ketamine: Teaching an old drug new tricks. *Anesth Analg* 1998; 87: pp. 1186-1193
[View In Article](#)
408. Kolodny A., Courtwright D.T., Hwang C.S., et al: The prescription opioid and heroin crisis: a public health approach to an epidemic of addiction. *Annu Rev Public Health* 2015; 36: pp. 559-574
[View In Article](#) | [Cross Ref](#)
409. Kolodny A., Courtwright D.T., Hwang C.S., Kreiner P., Eadie J.L., Clark T.W., et al: The prescription opioid and heroin crisis: a public health approach to an epidemic of addiction. *Annu Rev Public Health* 2015 Mar 18; 36: pp. 559-574
[View In Article](#) | [Cross Ref](#)
410. Koppert W., Ostermeier N., Sittl R., et al: Low-dose lidocaine reduces secondary hyperalgesia by a central mode of action. *Pain* 2000; 85: pp. 217-224
[View In Article](#) | [Cross Ref](#)
411. Kranke P., Jokinen J., Pace N.L., et al: Continuous intravenous perioperative lidocaine infusion for postoperative pain and recovery. *Cochrane Database Syst Rev* 2015; undefined:
[View In Article](#)

412. Krex D, Klink B, Hartmann C, von Deimling A, Pietsch T, Simon M, Sabel M, Steinbach JP, Heese O, Reifenberger G, Weller M, Schackert G (2007) Long-term survival with glioblastoma multiforme. *Brain* 130(Pt 10):2596–2606. <https://doi.org/10.1093/brain/awm204> CrossRef PubMed Google Scholar
413. Kumar S., et al: A comparative assessment of the clinical efficacy of intranasal desmopressin spray and diclofenac in the treatment of renal colic. *Urol Res* 2011; 39: pp. 397-400
[View In Article](#) | [Cross Ref](#)
414. Kurdi M.S., Theerth K.A., and Deva R.S.: Ketamine: current applications in anesthesia, pain, and critical care. *Anesth Essays Res* 2014; 8: pp. 283-289
[View In Article](#) | [Cross Ref](#)
415. Kurdi M.S., Theerth K.A., and Deva R.S.: Ketamine: current applications in anesthesia, pain, and critical care. *Anesth Essays Res* 2014; 8: pp. 283-289
[View In Article](#) | [Cross Ref](#)
416. Kurdi M.S., Theerth K.A., and Deva R.S.: Ketamine: Current applications in anesthesia, pain, and critical care. *Anesth Essays Res* 2014; 8: pp. 283-289
[View In Article](#) | [Cross Ref](#)
417. Kurdi MS, Theerth KA, Deva RS. Ketamine: Current applications in anesthesia, pain, and critical care. *Anesth Essays Res.* 2014;8(3):283–9. [PMC free article] [PubMed]
418. Kurdi MS, Theerth KA, Deva RS. Ketamine:Current applications in anesthesia, pain, and critical care. *Anesth Essays Res.* 2014;8(3):283–9. [PMC free article] [PubMed]
419. Lada P, Delgado G Jr. Documentation of pharmacists' interventions in an emergency department and associated cost avoidance. *Am J Health-Syst Pharm.* 2007; 64:63-8.
420. Laffel G., and Blumenthal D.: The case for using industrial quality management science in health care organizations. *JAMA* 1989; 262: pp. 2869-2873
[View In Article](#) | [Cross Ref](#)
421. LaFond C.M., Van Hulle Vincent C., Corte C., et al: PICU nurses' pain assessments and intervention choices for virtual human and written vignettes. *J Pediatr Nurs* 2015; 30: pp. 580-590
[View In Article](#) | [Cross Ref](#)
422. Laivenieks N, McCaul K, O'Brodovich M. Clinical pharmacy services provided to an emergency department. *Can J Hosp Pharm.* 1992; 45:113-5.
423. Lauretti G.R.: Mechanisms of analgesia of intravenous lidocaine. *Rev Bras Anestesiol* 2008; 58: pp. 280-286
[View In Article](#) | [Cross Ref](#)
424. Leape LL, Brennan TA, Laird N et al. The nature of adverse events in hospitalized patients. Results of the Harvard Medical Practice Study II. *N Engl J Med.* 1991; 324:377-84.
425. Leape LL. Pharmacist participation on physician rounds and adverse drug events in the intensive care unit. *JAMA.* 1999; 282:267-70. [Erratum, *JAMA.* 2000; 283: 1293.]
426. Lee J. Association between patient race/ethnicity and perceived interpersonal aspects of care in the ED. University of North Carolina. SAEM Abstracts. *Ann Emerg Med.* 2006;48:S121.

427. Lee T.H.: Zero pain is not the goal. *JAMA* 2016 Apr 19; 315: pp. 1575-1577
[View In Article](#) | [Cross Ref](#)
428. Lee T.S., and Hou X.: Vasoactive effects of ketamine on isolated rabbit pulmonary arteries. *Chest* 1995; 107: pp. 1152-1155
[View In Article](#) | [Cross Ref](#)
429. Lembke A. Why doctors prescribe opioids to known opioid abusers. *N Engl J Med.* 2012; 367:1580-1.
430. Lester L, Braude DA, Niles C, Crandall CS. Low-dose ketamine for analgesia in the ED:a retrospective case series. *Am J Emerg Med.* 2010;28(7):820-7. [[PubMed](#)]
431. Lester L, Braude DA, Niles C, Crandall CS. Low-dose ketamine for analgesia in the ED: a retrospective case series. *Am J Emerg Med* 2010;28:820-7.
432. Lester L, Braude DA, Niles C, et al. Low-dose ketamine for analgesia in the ED: a retrospective case series. *Am J Emerg Med.* 2010;28(7):820-7. [[PubMed](#)]
433. Lester L., Braude D.A., Niles C., and Crandall C.S.: Low-dose ketamine for analgesia in the ED: a retrospective case series. *Am J Emerg Med* 2010; 28: pp. 820-876
[View In Article](#) | [Cross Ref](#)
434. Lester L., Braude D.A., Niles C., et al: Low-dose ketamine for analgesia in the ED: a retrospective case series. *Am J Emerg Med* 2010; 28: pp. 820-827
[View In Article](#) | [Cross Ref](#)
435. Lester L.M., Braude D.M., Niles C.M., et al: Low-dose ketamine for analgesia in the ED: a retrospective case series. *Am J Emerg Med* 2010; 28: pp. 820-827
[View In Article](#) | [Cross Ref](#)
436. Lev R., Lee O., Petro S., et al: Who is prescribing controlled medications to patients who die of prescription drug abuse? *Am J Emerg Med* 2016; 34: pp. 30-35
[View In Article](#) | [Cross Ref](#)
437. Levy DB. Documentation of clinical and cost-saving pharmacy interventions in the emergency room. *Hosp Pharm.* 1993; 28:624-53.
438. Lewis LM, Lasater LC, Brooks CB. Are emergency physicians too stingy with analgesics? *South Med J.* 1994;87:7-9. [[PubMed](#)]
439. Lewis LM, Lasater LC, Ruoff BE. Failure of a chest pain clinical policy to modify physician evaluation and management. *Ann Emerg Med* 1995; 25:9-14.
440. Lewis LM, Lasater LC, Ruoff BE. Failure of a chest pain clinical policy to modify physician evaluation and management. *Ann Emerg Med.* 1995;25:9-14.
[[PubMed](#)]
441. Li K., Harris B., Hadi S., et al: What should be the optimal cut points for mild, moderate, and severe pain? *J Palliat Med* 2007; 10: pp. 1338-1346
[View In Article](#) | [Cross Ref](#)
442. Li K., Harris B., Hadi S., et al: What should be the optimal cut points for mild, moderate, and severe pain? *J Palliat Med* 2007; 10: pp. 1338-1346
[View In Article](#)
443. Liberati A., Altman D.G., Tetzlaff J., et al: The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med* 2009; 6: pp. e1000100
[View In Article](#) | [Cross Ref](#)

444. Liebmann O., Price D., Mills C., et al: Feasibility of forearm ultrasonography-guided nerve blocks of the radial, ulnar, and median nerves for hand procedures in the emergency department. *Ann Emerg Med* 2006; 48: pp. 558-562
[View In Article](#) | [Cross Ref](#)
445. Lin P.L., Fan S.Z., Huang C.H., et al: Analgesic effect of lidocaine patch 5% in the treatment of acute herpes zoster: a double-blind and vehicle-controlled study. *Reg Anesth Pain Med* 2008; 33: pp. 320-325
[View In Article](#)
446. Ling JM, Mike LA, Rubin J et al. Documentation of pharmacist interventions in the emergency department. *Am J HealthSyst Pharm.* 2005; 62:1793-7.
447. Linnebur S.A., O'Connell M.B., Wessell A.M., et al: Pharmacy practice, research, education, and advocacy for older adults. ACCP white paper. *Pharmacotherapy* 2005; 25: pp. 1404-1405
[View In Article](#)
448. Long J., and Shenvi C.: Acute pain and procedural sedation in older adults. In Strayer R., Motov S., and Nelson L. (eds): *Management of pain and procedural sedation in acute care*.
[View In Article](#)
449. Lopes T., et al: An assessment of the clinical efficacy of intranasal desmopressin spray in the treatment of renal colic. *BJU Int* 2001; 87: pp. 322-325
[View In Article](#) | [Cross Ref](#)
450. Louis DN, Perry A, Reifenberger G, von Deimling A, Figarella-Branger D, Cavenee WK, Ohgaki H, Wiestler OD, Kleihues P, Ellison DW (2016) The 2016 World Health Organization classification of tumors of the central nervous system: a summary. *Acta Neuropathol* 131(6):803–820. <https://doi.org/10.1007/s00401-016-1545-1> [CrossRef](#)[PubMed](#)[Google Scholar](#)
451. Luhmann J.D., Schootman M., Luhmann S.J., et al: A randomized comparison of nitrous oxide plus hematoma block versus ketamine plus midazolam for emergency department forearm fracture reduction in children. *Pediatrics* 2006; 118: pp. e1078-e1086
[View In Article](#) | [Cross Ref](#)
452. Luhmann J.D., Schootman M., Luhmann S.J., et al: A randomized comparison of nitrous oxide plus hematoma block versus ketamine plus midazolam for emergency department forearm fracture reduction in children. *Pediatrics* 2006; 118: pp. e1078-e1086
[View In Article](#) | [Cross Ref](#)
453. Lvovschi V., Auburn F., Bonnet P., et al: Intravenous morphine titration to treat severe pain in the ED. *Am J Emerg Med* 2008; 26: pp. 676-682
[View In Article](#) | [Cross Ref](#)
454. Lvovschi V., Auburn F., Bonnet P., et al: Intravenous morphine titration to treat severe pain in the ED. *Am J Emerg Med* 2008; 26: pp. 676-682
[View In Article](#) | [Cross Ref](#)
455. Macario A, Royal MA. A literature review of randomized clinical trials of intravenous acetaminophen (paracetamol) for acute postoperative pain. *Pain Pract* 2011;11:290–6.

456. MacDonald N., and MacLeod S.M.: Has the time come to phase out codeine? CMAJ 2010 Nov 23; 182: pp. 1825
[View In Article](#) | [Cross Ref](#)
457. Macintyre P.E.: Safety and efficacy of patient-controlled analgesia. Br J Anaesth 2001; 87: pp. 36-46
[View In Article](#) | [Cross Ref](#)
458. Maimonides Medical Center. Presenter Evaluation Form. Available at: http://cmetracker.net/MMC/evaluation_template.doc. Accessed Feb 20, 2011.
459. Majidinejad S, Esmailian M, Emadi M. Comparison of intravenous ketamine with morphine in pain relief of long bones fractures:a double blind randomized clinical trial. Emerg (Tehran) 2014;2(2):77–80. [[PMC free article](#)] [[PubMed](#)]
460. Malhotra AK, Pinals DA, Weingartner H, et al. NMDA receptor function and human cognition: the effects of ketamine in healthy volunteers. *Neuropsychopharmacology* 1996;14:301-7.
461. Manchikanti L., Helm S., Fellows B., et al: Opioid epidemic in the United States. Pain Physician 2012; 15: pp. ES9-38
[View In Article](#)
462. Mao J., and Chen L.L.: Systemic lidocaine for neuropathic pain relief. Pain 2000; 87: pp. 7-17
[View In Article](#) | [Cross Ref](#)
463. Marchertiene 2008.
[View In Article](#)
464. Marchertiene 2008.
[View In Article](#)
465. Marchertiene 2008.
[View In Article](#)
466. Marchertiene 2008.
[View In Article](#)
467. Marchertiene 2008.
[View In Article](#)
468. Marchertienè 2008.
[View In Article](#)
469. Marchertiene I.A., Macas A., and Karbonskiene A.: Pulmonary edema and hemorrhage as complications of acute airway obstruction following anesthesia. Medicina 2008; 44: pp. 871-876
[View In Article](#)
470. Marquié L, Raufaste E, Lauque D, Mariné C, Ecoiffier M, Sorum P. Pain rating by patients and physicians: evidence of systematic pain miscalibration. *Pain*. 2003; 102:289-96.
471. Marquié L, Raufaste E, Lauque D, Mariné C, Ecoiffier M, Sorum P. Pain rating by patients and physicians: Evidence of systematic pain miscalibration. *Pain*. 2003;102:289–296. [[PubMed](#)]
472. Marret E., Rolin M., Beaussier M., et al: Meta-analysis of intravenous lidocaine and postoperative recovery after abdominal surgery. Br J Surg 2008; 95: pp. 1331-1338
[View In Article](#) | [Cross Ref](#)

473. Masoumi K, Forouzan A, Asgari Darian A, Feli M, Barzegari H, Khavanin A. Comparison of clinical efficacy of intravenous acetaminophen with intravenous morphine in acute renal colic: a randomized, double-blind, controlled trial. *Emerg Med Int* 2014;2014:571326.
474. Masoumi K, Forouzan A, Darian AA et al. Comparison of clinical efficacy of intravenous acetaminophen with intravenous morphine in acute renal colic: a randomized, double-blind, controlled trial. *Emerg Med Int*. 2014; 2014:571326.
475. Masoumi K., et al: The efficacy of intranasal desmopressin as an adjuvant in the acute renal colic pain management. *Pain Res Treat* 2014; 2014: [View In Article](#)
476. Masoumi K., Forouzan A., Asgari Darian A., et al: Comparison of clinical efficacy of intravenous acetaminophen with intravenous morphine in acute renal colic: a randomized, double-blind, controlled trial. *Emerg Med Int* 2014; 2014: pp. 571326 [View In Article](#)
477. Mazer-Amirshahi M, Mullins PM, Rasooly I et al. Rising opioid prescribing in adult US emergency department visits: 2001-2010. *Acad Emerg Med*. 2014; 21:236-43.
478. McCarberg B., and D'Arcy Y.: Options in topical therapies in the management of patients with acute pain. *Postgrad Med* 2013; 125: pp. 19-24 [View In Article](#) | [Cross Ref](#)
479. McCarthy G.C., Megalla S.A., and Habib A.S.: Impact of intravenous lidocaine infusion on postoperative analgesia and recovery from surgery: a systematic review of randomized controlled trials. *Drugs* 2010; 70: pp. 1149-1163 [View In Article](#) | [Cross Ref](#)
480. McCleane G.: Intravenous lidocaine: an outdated or underutilized treatment for pain? *J Palliat Med* 2007; 10: pp. 798-805 [View In Article](#) | [Cross Ref](#)
481. McGhie J., and Serpell M.G.: Clinical pharmacology: local anesthetics. In Macintyre P.E., Walker S.M., and Rowbotham D.J. (eds): Clinical pain management (acute pain), 2nd edition. London: Hodder & Stoughton Limited, 2008. pp. 113-129 [View In Article](#)
482. McGhie J., and Serpell M.G.: Clinical pharmacology: local anesthetics. In Macintyre P.E., Walker S.M., and Rowbotham D.J. (eds): Clinical pain management (acute pain), 2nd edition. London: Hodder & Stoughton Limited, 2008. pp. 113-129 [View In Article](#)
483. McMullin ST, Hennenfent JA, Ritchie DJ et al. A prospective, randomized trial to assess the cost impact of pharmacist-initiated interventions. *Arch Intern Med*. 1999; 159:2306-9.
484. McQuay H.: Opioids in pain management. *Lancet* 1999 Jun 26; 353: pp. 2229-2232 [View In Article](#) | [Cross Ref](#)
485. McReynolds T.M., and Sheridan B.J.: Intramuscular ketorolac versus osteopathic manipulative treatment in the management of acute neck pain in the emergency department: a randomized clinical trial. *J Am Osteopath Assoc* 2005; 105: pp. 57-68 [View In Article](#)
486. Melendez 2009. [View In Article](#)

487. Melendez E., and Bachur R.: Serious adverse events during procedural sedation with ketamine. *Pediatr Emerg Care* 2009; 25: pp. 325-328
[View In Article](#) | [Cross Ref](#)
488. Mercadante S, Lodi F, Sapiro M, Calligara M, Serretta R. Long-term ketamine subcutaneous continuous infusion in neuropathic cancer pain. *J Pain Symptom Manage* 1995;10:564-8.
489. Meredith J.T., Wait S., and Brewer K.L.: A prospective double-blind study of nasal sumatriptan versus IV ketorolac in migraine. *Am J Emerg Med* 2003; 21: pp. 173-175
[View In Article](#) | [Cross Ref](#)
490. Messenger DW, Murray HE, Dungey PE, van Vlymen J, Sivilotti. Subdissociative-dose ketamine versus fentanyl for analgesia during propofol procedural sedation: a randomized clinical trial. *Acad Emerg Med* 2008;15:877-86.
491. Mezei L., and Murinson B.B.: Pain education in North American medical schools. *J Pain* 2011; 12: pp. 1199-1208
[View In Article](#) | [Cross Ref](#)
492. Mialon PJ, Williams P, Wiebe RA. Clinical pharmacy services in a pediatric emergency room. *Hosp Pharm*. 2004; 39:121-4.
493. Miller A.C., Jamin C.T., and Elamin E.M.: Continuous intravenous infusion of ketamine for maintenance sedation. *Minerva Anesth* 2011; 77: pp. 812-820
[View In Article](#)
494. Miller G.E.: The assessment of clinical skills/competence/performance. *Acad Med J Assoc Am Med Coll* 1990; 65: pp. S63-S67
[View In Article](#)
495. Miller J., Schauer S., Ganem V., et al: Low-dose ketamine vs morphine for acute pain in the ED: a randomized controlled trial. *Am J Emerg Med* 2015; undefined:
[View In Article](#)
496. Miller J., Schauer S., Ganem V., et al: Low-dose ketamine vs morphine for acute pain in the ED: a randomized controlled trial. *Am J Emerg Med* 2015; 33: pp. 402-408
[View In Article](#) | [Cross Ref](#)
497. Miller J.P., Schauer S.G., Ganem V.J., and Bebarta V.S.: Low-dose ketamine vs morphine for acute pain in the ED: a randomized controlled trial. *Am J Emerg Med* 2015 Mar; 33: pp. 402-408
[View In Article](#) | [Cross Ref](#)
498. Miller J.P., Schauer S.G., Ganem V.J., and Bebarta V.S.: Low-dose ketamine vs morphine for acute pain in the ED: a randomized controlled trial. *Am J Emerg Med* 2015; 33: pp. 402-408
[View In Article](#) | [Cross Ref](#)
499. Miller J.P., Schauer S.G., Ganem V.J., et al: Low-dose ketamine vs morphine for acute pain in the ED: a randomized controlled trial. *Am J Emerg Med* 2015; 33: pp. 402-408
[View In Article](#) | [Cross Ref](#)
500. Miller JP, Schauer SG, Ganem VJ. Lowdose ketamine vs morphine for acute pain in the ED: a randomized controlled trial. *Am J Emerg Med*. 2015; 33:402-8.
501. Miller M., Barber C.W., Leatherman S., Fonda J., Hermos J.A., Cho K., et al: Prescription opioid duration of action and the risk of unintentional overdose among

patients receiving opioid therapy. *JAMA Intern Med* 2015 Apr; 175: pp. 608-615
[View In Article](#) | [Cross Ref](#)

502. Miner J.R., Kletti C., Herold M., et al: Randomized clinical trial of nebulized fentanyl citrate versus i.v. fentanyl citrate in children presenting to the emergency department with acute pain. *Acad Emerg Med* 2007; 14: pp. 895-898
[View In Article](#)
503. Miner J.R., Kletti C., Herold M., et al: Randomized clinical trial of nebulized fentanyl citrate versus i.v. fentanyl citrate in children presenting to the emergency department with acute pain. *Acad Emerg Med* 2007; 14: pp. 895-898
[View In Article](#)
504. Minotti V., Betti M., Ciccarese G., Fumi G., Tonato M., Del Favero A. A double-blind study comparing two single-dose regimens of ketorolac with diclofenac in pain due to cancer. *Pharmacotherapy*. 1998;18(3):504–8. [[PubMed](#)]
505. Minotti V., Betti M., Ciccarese G., et al: A double-blind study comparing two single-dose regimens of ketorolac with diclofenac in pain due to cancer. *Pharmacotherapy* 1998; 18: pp. 504-508
[View In Article](#)
506. Minotti V., Betti M., Ciccarese G., et al: A double-blind study comparing two single-dose regimens of ketorolac with diclofenac in pain due to cancer. *Pharmacotherapy* 1998; 18: pp. 504-508
[View In Article](#)
507. Mitchell A., and Fallon M.: A single infusion of intravenous ketamine improves pain relief in patients with critical limb ischaemia: results of a double blind randomised controlled trial. *Pain* 2002; 97: pp. 275-281
[View In Article](#) | [Cross Ref](#)
508. Mizoe JE, Tsujii H, Hasegawa A, Yanagi T, Takagi R, Kamada T, Tsuji H, Takakura K (2007) Phase I/II clinical trial of carbon ion radiotherapy for malignant gliomas: combined X-ray radiotherapy, chemotherapy, and carbon ion radiotherapy. *Int J Radiat Oncol Biol Phys* 69(2):390–396. <https://doi.org/10.1016/j.ijrobp.2007.03.003> [CrossRef](#)[PubMed](#)[Google Scholar](#)
509. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 2009;62:1006–12.
510. Moher D., Liberati A., Tetzlaff J., Altman D.G., and PRISMA Group : Preferred reporting items for systematic reviews and meta-analysis: the PRISMA statement. *J Clin Epidemiol* 2009; 62: pp. 1006-1012
[View In Article](#) | [Cross Ref](#)
511. Montori V.M., Swiontkowski M.F., and Cook D.J.: Methodologic issues in systematic reviews and meta-analyses. *Clin Orthop Relat Res* 2003; 413: pp. 43-54
[View In Article](#) | [Cross Ref](#)
512. Moore P.A., and Hersh E.V.: Combining ibuprofen and acetaminophen for acute pain management after third-molar extractions: translating clinical research to dental practice. *J Am Dent Assoc* 2013 Aug; 144: pp. 898-908
[View In Article](#) | [Cross Ref](#)
513. Morley-Forster P.K., Pergolizzi J.V., Taylor R., Axford-Gatley R.A., and Sellers E.M.: Mitigating the risk of opioid abuse through a balanced undergraduate pain

medicine curriculum. *J Pain Res* 2013; 6: pp. 791-801

[View In Article](#) | [Cross Ref](#)

514. Morone N.E., and Weiner D.K.: Pain as the fifth vital sign: exposing the vital need for pain education. *Clin Ther* 2013; 35: pp. 1728-1732
[View In Article](#) | [Cross Ref](#)
515. Morteza-Bagi HR, Amjadi M, Mirzaii-Sousefidi R. The comparison of Apotel plus low dose morphine and full dose of morphine in pain relief in patients with acute renal colic. *Addict Health* 2015;7:66-73.
516. Morton N.S.: Ketamine for procedural sedation and analgesia in pediatric emergency medicine: a UK perspective. *Paediatr Anaesth* 2008; 18: pp. 25-29
[View In Article](#) | [Cross Ref](#)
517. Mosier J, Roper G, Hays D et al. Sedative dosing or propofol for treatment of migraine headache in the emergency department: a case series. *West J Emerg Med.* 2013; 14:646-9.
518. Motov S, Drapkin J, Likourezos A, Beals T, Monfort R, Fromm C, et al. Continuous intravenous sub-dissociative dose ketamine infusion for managing pain in the emergency department. *West J Emerg Med.* 2018;19(3):559–66. [PMC free article] [\[PubMed\]](#)
519. Motov S, Mai M, Pushkar I, et al. A prospective randomized, double-dummy trial comparing intravenous push dose of low dose ketamine to short infusion of low dose ketamine for treatment of moderate to severe pain in the emergency department. *Am J Emerg Med.* 2017;35(8):1095–100. [\[PubMed\]](#)
520. Motov S, Mai M, Pushkar I, Likourezos A, Drapkin J, Yasavolian M, et al. A prospective randomized, double-dummy trial comparing IV push low dose ketamine to short infusion of low dose ketamine for treatment of pain in the ED. *Am J Emerg Med.* 2017;35(8):1095–100. [\[PubMed\]](#)
521. Motov S, Rockoff B, Cohen V et al. Intravenous subdissociative-dose ketamine versus morphine for analgesia in the emergency department: a randomized controlled trial. *Ann Emerg Med.* 2015; 66:222–9.e1.
522. Motov S, Rockoff B, Cohen V, et al. Intravenous subdissociative-dose ketamine versus morphine for analgesia in the emergency department: a randomized controlled trial. *Ann Emerg Med.* 2015;66(3):222–9. [\[PubMed\]](#)
523. Motov S, Rockoff B, Cohen V, Pushkar I, Likourezos A, McKay C, et al. Intravenous subdissociative-dose ketamine versus morphine for analgesia in the emergency department:a randomized controlled trial. *Ann Emerg Med.* 2015;66(3):222–9. [\[PubMed\]](#)
524. Motov S, Strayer R, Hayes B, et al. AAEM White Paper on acute pain management in the emergency department. American Academy of Emergency Medicine. Available at: <http://www.aaem.org/em-resources/position-statements/2017/pain-mgmt-in-ed>. Accessed December 1, 2017.
[View In Article](#)
525. Motov S., Mai M., Pushkar I., et al: A prospective randomized, double-dummy trial comparing intravenous push dose of low dose ketamine to short infusion of low dose ketamine for treatment of moderate to severe pain in the emergency department. *Am J Emerg Med* 2017; 35: pp. 1095-1100
[View In Article](#) | [Cross Ref](#)

526. Motov S., Mai M., Pushkar I., et al: A prospective randomized, double-dummy trial comparing IV push low dose ketamine to short infusion of low dose ketamine for treatment pain in the ED. *Am J Emerg Med* 2017; 35: pp. 1095-1100
[View In Article](#) | [Cross Ref](#)
527. Motov S., Rockoff B., Cohen V., et al: Intravenous subdissociative-dose ketamine versus morphine for analgesia in the emergency department: a randomized controlled trial. *Ann Emerg Med* 2015; 66: pp. 222-229
[View In Article](#) | [Cross Ref](#)
528. Motov S., Rockoff B., Cohen V., et al: Intravenous subdissociative-dose ketamine versus morphine for analgesia in the emergency department: a randomized controlled trial. *Ann Emerg Med* 2015; 66: pp. 222-229
[View In Article](#) | [Cross Ref](#)
529. Motov S., Rockoff B., Cohen V., et al: Intravenous subdissociative-dose ketamine versus morphine for analgesia in the emergency department: a randomized controlled trial. *Ann Emerg Med* 2015; 66: pp. 222-229
[View In Article](#) | [Cross Ref](#)
530. Motov S., Rockoff B., Cohen V., et al: Intravenous subdissociative-dose ketamine versus morphine for analgesia in the emergency department: a randomized controlled trial. *Ann Emerg Med* 2015; 66: pp. 222-229.e1
[View In Article](#) | [Cross Ref](#)
531. Motov S., Rockoff B., Cohen V., Pushkar I., Likourezos A., McKay C., et al: Intravenous sub-dissociative-dose ketamine versus morphine for analgesia in the emergency department: a randomized controlled trial. *Ann Emerg Med* 2015; 22: pp. 251-257
[View In Article](#) | [Cross Ref](#)
532. Motov S., Rockoff B., Cohen V., Pushkar I., Likourezos A., McKay C., et al: Intravenous subdissociative-dose ketamine versus morphine for analgesia in the emergency department: a randomized controlled trial. *Ann Emerg Med* 2015 Sep; 66: pp. 222-229.e1
[View In Article](#) | [Cross Ref](#)
533. Motov S.M., and Marshall J.P.: Acute pain management curriculum for emergency medicine residency programs. *Acad Emerg Med* 2011; 18: pp. S87-S91
[View In Article](#) | [Cross Ref](#)
534. Motov S.M., and Nelson L.S.: Advanced concepts and controversies in emergency department pain management. *Anesthesiol Clin* 2016; 34: pp. 271-285
[View In Article](#) | [Cross Ref](#)
535. Motov S.M., and Nelson L.S.: Advanced concepts and controversies in emergency department pain management. *Anesthesiol Clin* 2016; 34: pp. 271-285
[View In Article](#) | [Cross Ref](#)
536. Motov SM, Khan AN. Problems and barriers of pain management in the emergency department: are we ever going to get better? *J Pain Res*. 2009; 2:5-11.
537. Motov SM, Marshall JP. Acute pain management curriculum for emergency medicine residency programs: acute pain management curriculum for the EM resident. *Acad Emerg Med*. 2011; 18:S87-91.
538. Mroszczak E.J., Jung D., Yee J., et al: Ketorolac tromethamine pharmacokinetics and metabolism after intravenous, intramuscular, and oral .

Pharmacotherapy 1990; 10: pp. 33S-39S

[View In Article](#)

539. Murad M.H., Mustafa R.A., Schunemann H.J., et al: Rating the certainty in evidence in the absence of a single estimate of effect. Evid Based Med 2017; 22: pp. 85-87
[View In Article](#) | [Cross Ref](#)
540. Murinson B.B., Gordin V., Flynn S., et al: Recommendations for a new curriculum in pain medicine for medical students: toward a career distinguished by competence and compassion. Pain Med 2013; 14: pp. 345-350
[View In Article](#) | [Cross Ref](#)
541. Murphy J.L.: Hypertension and pulmonary edema associated with ketamine administration in a patient with a history of substance abuse. Can J Anesth 1993; 40: pp. 160-164
[View In Article](#) | [Cross Ref](#)
542. Nakayama M., and Murray P.A.: Ketamine preserves and propofol potentiates hypoxic pulmonary vasoconstriction compared with the conscious state in chronically instrumented dogs. Anesthesiology 1999; 91: pp. 760
[View In Article](#) | [Cross Ref](#)
543. Nasca T.J., Philibert I., Brigham T., and Flynn T.C.: The next GME accreditation system—rationale and benefits. N Engl J Med 2012; 366: pp. 1051-1056
[View In Article](#) | [Cross Ref](#)
544. Neighbor M.L., and Puntillo K.A.: Intramuscular ketorolac vs oral ibuprofen in emergency department patients with acute pain. Acad Emerg Med 1998; 5: pp. 118-122
[View In Article](#) | [Cross Ref](#)
545. Neighbor ML, Honner S, Kohn MA. Factors affecting emergency department opioid administration to severely injured patients. *Acad Emerg Med.* 2004; 11:1290-6.
546. Neighbor ML, Honner S, Kohn MA. Factors affecting emergency department opioid administration to severely injured patients. *Acad Emerg Med.* 2004;11:1290–1296. [PubMed]
547. Nelson BP, Cohen D, Lander O, Crawford N, Viccellio AW, Singer AJ. Mandated pain scales improve frequency of ED analgesic administration. *Am J Emerg Med.* 2004;22:582–585. [PubMed]
548. Nelson L.S., and Juurlink D.N.: Tramadol and hypoglycemia: one more thing to worry about. JAMA Intern Med 2015 Feb; 175: pp. 194-195
[View In Article](#)
549. Nelson L.S., Juurlink D.N., and Perrone J.: Addressing the opioid epidemic. JAMA 2015 Oct 13; 314: pp. 1453-1454
[View In Article](#) | [Cross Ref](#)
550. Nesbit T, Shermock K, Bobek MB et al. Implementation and pharmacoeconomic analysis of a clinical staff pharmacist practice model. Am J Health-Syst Pharm. 2001; 58:784-90.
551. Neven D.E., Sabel J.C., Howell D.N., and Carlisle R.J.: The development of the Washington State emergency department opioid prescribing guidelines. J Med Toxicol Off J Am Coll Med Toxicol 2012; 8: pp. 353-359
[View In Article](#) | [Cross Ref](#)
552. Newton 2008.
[View In Article](#)

553. Newton 2008.
[View In Article](#)
554. Newton A., and Fitton L.: Intravenous ketamine for adult procedural sedation in the emergency department: A prospective cohort study. *Emerg Med J* 2008; 25: pp. 498
[View In Article](#) | [Cross Ref](#)
555. Nicol MF, Ashton-Cleary D. "Why haven't you taken any pain killers?" A patient focused study of the walking wounded in an urban emergency department. *Emerg Med J*. 2003;20:228–229. [\[PMC free article\]](#) [\[PubMed\]](#)
556. Niesters M, Martini C, Dahan A. Ketamine for chronic pain: risks and benefits. *Br J Clin Pharmacol* 2014;77:357–67.
557. Nisbet A.C.: Intramuscular gluteal injections in the increasing obese population: retrospective study. *BMJ* 2006; 332: pp. 637-638
[View In Article](#) | [Cross Ref](#)
558. Nishimoto RN. OFIRMEV: an old drug becomes new again. *Anesth Prog* 2014;61:99–102.
559. Nursing Care Quality Assurance Commission. Administration of sedating, analgesic and anesthetic agents. [www. doh.wa.gov/Portals/1/Documents/6000/AdminOfSedating,AnalgesicAndAnesthetic Agents.pdf](http://www.doh.wa.gov/Portals/1/Documents/6000/AdminOfSedating,AnalgesicAndAnesthetic%20Agents.pdf) (accessed 2015 May 13).
560. O'Donnell J, Ferguson LP, Beattie TF. Use of analgesia in a paediatric accident and emergency department following limb trauma. *Eur J Emerg Med*. 2002;9:5–8. [\[PubMed\]](#)
561. O'Connor A., Schug S.A., and Cardwell H.: A comparison of the efficacy and safety of morphine and pethidine as analgesia for suspected renal colic in the emergency setting. *J Accid Emerg Med* 2000; 17: pp. 261-264
[View In Article](#) | [Cross Ref](#)
562. Office of National Drug Control Policy. Epidemic: responding to America's prescription drug abuse crisis. [www. whitehouse.gov/sites/default/files/ondcp/ policy-and-research/rx_abuse_plan.pdf](http://www.whitehouse.gov/sites/default/files/ondcp/policy-and-research/rx_abuse_plan.pdf) (accessed 2014 Nov 1).
563. Ogata T, Teshima T, Inaoka M, Minami K, Tsuchiya T, Isono M, Furusawa Y, Matsuura N (2011) Carbon Ion Irradiation Suppresses Metastatic Potential of Human Non-small Cell Lung Cancer A549 Cells through the Phosphatidylinositol-3-Kinase/Akt Signaling Pathway. *J Radiat Res* 52(3):374–379. <https://doi.org/10.1269/jrr.10102> [CrossRef](#)[PubMed](#)[Google Scholar](#)
564. Oguzturk J, Ozgur D, Turtay MG, et al. Tramadol or paracetamol do not effect the diagnostic accuracy of acute abdominal pain with significant pain relief - a prospective, randomized, placebo controlled double blind study. *Eur Rev Med Pharmacol Sci* 2012;16:1983–8.
565. Olsen J.C., Ogarek J.L., Goldenberg E.J., and Sulo S.: Impact of a chronic pain protocol on emergency department utilization. *Acad Emerg Med* 2016 Apr; 23: pp. 424-432
[View In Article](#) | [Cross Ref](#)
566. Oneschuk D, Fainsinger R, Hanson J, Bruera E. Assessment and knowledge in palliative care in second year family medicine residents. *J Pain Symptom Manage*. 1997;14:265–273. [\[PubMed\]](#)
567. Ong C.K., Seymour R.A., Lirk P., et al: Combining paracetamol (acetaminophen) with nonsteroidal antiinflammatory drugs: a qualitative systematic review of analgesic

- efficacy for acute postoperative pain. Anesth Analg 2010; 110: pp. 1170-1179
[View In Article](#)
568. Oregon Chapter of American College of Emergency Physicians. Oregon emergency department (ED) opioid prescribing guidelines. www.ocep.org/images/pdf/ed_opioid_abuse_guidelines.pdf (accessed 2014 Nov 1).
569. O'Sullivan I., and Benger J.: Nitrous oxide in emergency medicine. Emerg Med J 2003; 20: pp. 214-217
[View In Article](#) | [Cross Ref](#)
570. Øye I, Paulsen O, Maursset A. Effects of ketamine on sensory perception: evidence for a role of N-methyl-D-aspartate receptors. *J Pharmacol Exp Ther* 1992;260:1209-13.
571. Painless ER website. Available at: www.painfree-ed.com. Accessed November 28, 2015.
[View In Article](#)
572. Pandey 2000.
[View In Article](#)
573. Pandey C.K., Mathur N., Singh N., and Chandola H.C.: Fulminant pulmonary edema after intramuscular ketamine. Can J Anesth. 2000; 47: pp. 894-896
[View In Article](#) | [Cross Ref](#)
574. Paqueron X., Lumbroso A., Mergoni P., et al: Is morphine-induced sedation synonymous with analgesia during intravenous morphine titration? Br J Anaesth 2002; 89: pp. 697-701
[View In Article](#)
575. Park CM, Park MJ, Kwak HJ, Lee HC, Kim MS, Lee SH, Park IC, Rhee CH, Hong SI (2006) Ionizing radiation enhances matrix metalloproteinase-2 secretion and invasion of glioma cells through Src/epidermal growth factor receptor-mediated p38/Akt and phosphatidylinositol 3-kinase/Akt signaling pathways. *Cancer Res* 66(17):8511-8519. <https://doi.org/10.1158/0008-5472.can-05-4340> CrossRefPubMedGoogle Scholar
576. Parthasarathy 2009.
[View In Article](#)
577. Parthasarathy 2009.
[View In Article](#)
578. Parthasarathy S., Ravishankar M., Selvarajan S., and Anbalagan T.: Ketamine and pulmonary edema-- Report of two cases. Indian J Anesth 2009; 53: pp. 486-488
[View In Article](#)
579. Patanwala A.E., Edwards C.J., Stoltz L., et al: Should morphine dosing be weight based for analgesia in the emergency department? J Opioid Manag 2012; 8: pp. 51-55
[View In Article](#) | [Cross Ref](#)
580. Patanwala A.E., Edwards C.J., Stoltz L., et al: Should morphine dosing be weight based for analgesia in the emergency department? J Opioid Manag 2012; 8: pp. 51-55
[View In Article](#)
581. Patanwala A.E., Holmes K.L., and Erstad B.L.: Analgesic response to morphine in obese and morbidly obese patients in the emergency department. Emerg Med J 2014; 31: pp. 139-142
[View In Article](#) | [Cross Ref](#)

582. Patanwala A.E., Keim S.M., and Erstad B.L.: Intravenous opioids for severe acute pain in the emergency department. Ann Pharmacother 2010; 44: pp. 1800-1809
[View In Article](#) | [Cross Ref](#)
583. Patanwala A.E., Keim S.M., and Erstad B.L.: Intravenous opioids for severe acute pain in the emergency department. Ann Pharmacother 2010; 44: pp. 1800-1809
[View In Article](#) | [Cross Ref](#)
584. Pathan S.A., Mitra B., Straney L.D., Afzal M.S., Anjum S., Shukla D., et al: Delivering safe and effective analgesia for management of renal colic in the emergency department: a double-blind, multigroup, randomised controlled trial. Lancet 2016 Mar; 15:
[View In Article](#)
585. Pathan S.A., Mitra B., Straney L.D., et al: Delivering safe and effective analgesia for management of renal colic in the emergency department: a double-blind, multigroup, randomised controlled trial. Lancet 2016; 387: pp. 1999-2007
[View In Article](#) | [Cross Ref](#)
586. Patil S., and Anitescu M.: Efficacy of outpatient ketamine infusions in refractory chronic pain syndromes: a 5-year retrospective analysis. 2012; 13: pp. 263-269
[View In Article](#) | [Cross Ref](#)
587. Paulozzi L.J.: Prescription drug overdoses: a review. J Safety Res 2012; 43: pp. 283-289
[View In Article](#) | [Cross Ref](#)
588. Paw I, Carpenter RC, Watabe K, Debinski W, Lo HW (2015) Mechanisms regulating glioma invasion. Cancer Lett 362(1):1–7. <https://doi.org/10.1016/j.canlet.2015.03.015>
[CrossRef](#)[PubMed](#)[PubMed Central](#)[Google Scholar](#)
589. Peirce R.J., Fragen R.J., and Pemberton D.M.: Intravenous ketorolac tromethamine versus morphine sulfate in the treatment of immediate postoperative pain. Pharmacotherapy 1990; 10: pp. 111s-115s
[View In Article](#)
590. Peirce RJ, Fragen RJ, Pemberton DM. Intravenous ketorolac tromethamine versus morphine sulfate in the treatment of immediate postoperative pain. Pharmacotherapy. 1990;10(6 Pt 2):111S–115S. [\[PubMed\]](#)
591. Peixoto R.D., and Hawley P.: Intravenous lidocaine for cancer pain without electrocardiographic monitoring: a retrospective review. J Palliat Med 2015; 18: pp. 373-377
[View In Article](#) | [Cross Ref](#)
592. Perrone J, DeRoos FJ, Nelson LS. Prescribing practices, knowledge, and use of prescription drug monitoring programs (PDMP) by a national sample of medical toxicologists, 2012. J Med Toxicol. 2012; 8:341-52.
593. Perrone J., Nelson L.S., and Yealy D.M.: Choosing analgesics wisely: what we know (and still need to know) about long-term consequences of opioids. Ann Emerg Med 2015; 65: pp. 500-502
[View In Article](#) | [Cross Ref](#)
594. Perrone J., Nelson L.S., and Yealy D.M.: Choosing analgesics wisely: what we know (and still need to know) about long-term consequences of opioids. Ann Emerg Med

2015; 65: pp. 500-502

[View In Article](#) | [Cross Ref](#)

595. Peth HA Jr. Medication errors in the emergency department: a systems approach to minimizing risk. *Emerg Med Clin North Am.* 2003; 21:141-58.
596. Petrenko AB, Yamakura T, Baba H, Shimoji K. The role of N-methyl-D-aspartate (NMDA) receptors in pain: a review. *Anesth Analg* 2003;97:1108-16.
597. Phillips D.M.: JCAHO pain management standards are unveiled. Joint Commission on Accreditation of Healthcare Organizations. *JAMA* 2000; 284: pp. 428-429
[View In Article](#)
598. Pickering G, Moustafa F, Macian N, Schmidt J, Pereira B, Dubray C. A new transmucous-buccal formulation of acetaminophen for acute traumatic pain: a non-inferiority, randomized, double-blind, clinical trial. *Pain Physician* 2015;18:249-57.
599. Pines JM, Hollander JE. Emergency department crowding is associate with poor care for patients with severe pain. *Ann Emerg Med.* 2008;51:1-5. [[PubMed](#)]
600. Pitetti R.D., Singh S., and Pierce M.C.: Safe and efficacious use of procedural sedation and analgesia by non-anesthesiologists in pediatric emergency department. *Arch Pediatr Adolesc Med* 2003; 157: pp. 1090.6
[View In Article](#)
601. Pizzo P.A., and Clark N.M.: Alleviating suffering 101—pain relief in the United States. *N Engl J Med* 2012; 366: pp. 197-199
[View In Article](#) | [Cross Ref](#)
602. Platts-Mills T.F., Esserman D.A., Brown D.L., Bortsov A.V., Sloane P.D., and McLean S.A.: Older US emergency department patients are less likely to receive pain medication than younger patients: results from a national survey. *Ann Emerg Med* 2012; 60: pp. 199-206
[View In Article](#) | [Cross Ref](#)
603. Platts-Mills T.F., Esserman D.A., Brown D.L., et al: Older US emergency department patients are less likely to receive pain medication than younger patients: results from a national survey. *Ann Emerg Med* 2012 Aug; 60: pp. 199-206
[View In Article](#) | [Cross Ref](#)
604. Pletcher M.J., Kertesz S.G., Kohn M.A., and Gonzales R.: Trends in opioid prescribing by race/ethnicity for patients seeking care in US emergency departments. *JAMA* 2008; 299: pp. 70-78
[View In Article](#)
605. Pletcher MJ, Kertesz SG, Kohn MA et al. Trends in opioid prescribing by race/ethnicity for patients seeking care in US emergency departments. *JAMA*. 2008; 299:70-8.
606. Pletcher MJ, Kertesz SG, Kohn MA, Gonzales R. Trends in opioid prescribing by race/ethnicity for patients seeking care in US emergency departments. *JAMA*. 2008;299:70-78. [[PubMed](#)]
607. Polivka J Jr, Polivka J, Holubec L, Kubikova T, Priban V, Hes O, Pivoarcikova K, Treskova I (2017) Advances in experimental targeted therapy and immunotherapy for patients with glioblastoma multiforme. *Anticancer Res* 37(1):21–33. <https://doi.org/10.21873/anticanres.11285>
[CrossRef](#)[PubMed](#)[Google Scholar](#)

608. Pollack C.V., Diercks D.B., Thomas S.H., Shapiro N.I., Fanikos J., Mace S.E., et al: Patient-reported outcomes from a national, prospective, observational study of emergency department acute pain management with an intranasal nonsteroidal anti-inflammatory drug, opioids, or both. *Acad Emerg Med* 2016 Mar; 23: pp. 331-341
[View In Article](#) | [Cross Ref](#)
609. Pomerleau A.C., Nelson L.S., Hoppe J.A., Salzman M., Weiss P.S., and Perrone J.: The impact of prescription drug monitoring programs and prescribing guidelines on emergency department opioid prescribing: a multi-center survey. *Pain Med* 2016 Mar; 19:
[View In Article](#)
610. Poon SJ, Greenwood-Erickson MB. The opioid prescription epidemic and the role of emergency medicine. *Ann Emerg Med*. 2014; 64:490-5.
611. Poonai N., Bhullar G., Lin K., Papini A., Mainprize D., Howard J., et al: Oral administration of morphine versus ibuprofen to manage postfracture pain in children: a randomized trial. *CMAJ*. 2014 Dec 9; 186: pp. 1358-1363
[View In Article](#) | [Cross Ref](#)
612. Portenoy R.K., and Foley K.M.: Chronic use of opioid analgesics in non-malignant pain: report of 38 cases. *Pain* 1986 May; 25: pp. 171-186
[View In Article](#) | [Cross Ref](#)
613. Porter J., and Jick H.: Addiction rare in patients treated with narcotics. *N Engl J Med* 1980 Jan 10; 302: pp. 123
[View In Article](#) | [Cross Ref](#)
614. Powell MF, Solomon DK, McEachen RA. Twenty-four hour emergency pharmaceutical services. *Am J Hosp Pharm*. 1985; 42:831-5.
615. Pribluda A, de la Cruz CC, Jackson EL (2015) Intratumoral heterogeneity: from diversity comes resistance. *Clin Cancer Res* 21(13):2916–2923. <https://doi.org/10.1158/1078-0432.ccr-14-1213> CrossRefPubMedGoogle Scholar
616. Raftery KA, Smith-Coggins R, Chen AH. Gender-associated differences in emergency department pain management. *Ann Emerg Med*. 1995;26:414–421. [PubMed]
617. Rahman N.H., and DeSilva T.: A randomized controlled trial of patient-controlled analgesia compared with boluses of analgesia for the control of acute traumatic pain in the emergency department. *J Emerg Med* 2012; 43: pp. 951-957
[View In Article](#) | [Cross Ref](#)
618. Ramasubbu C., and Gupta A.: Pharmacological treatment of opioid-induced hyperalgesia: a review of the evidence. *J Pain Palliat Care Pharmacother* 2011; 25: pp. 219-230
[View In Article](#) | [Cross Ref](#)
619. Rawlins JM, Khan AA, Shenton AF, Sharpe DT. Epidemiology and outcome analysis of 208 children with burns attending an emergency department. *Pediatr Emerg Care*. 2007;23:289–233. [PubMed]
620. Raychaudhuri B, Vogelbaum MA (2011) IL-8 is a mediator of NF-kappaB induced invasion by gliomas. *J Neuro-oncol* 101(2):227–235. <https://doi.org/10.1007/s11060-010-0261-2> CrossRefGoogle Scholar
621. Reifler LM, Droz D, Bailey JE et al. Do prescription monitoring programs impact state trends in opioid abuse/misuse? *Pain Med*. 2012; 13:434-42.

622. Reuben S.S., Connelly N.R., Lurie S., et al: Dose-response of ketorolac as an adjunct to patient-controlled analgesia morphine in patients after spinal fusion surgery. *Anesth Analg* 1998; 87: pp. 98-102
[View In Article](#)
623. Reutens D.C., Fatovich D.M., Stewart-Wynne E.G., et al: Is intravenous lidocaine clinically effective in acute migraine? *Cephalgia* 1991; 11: pp. 245-247
[View In Article](#) | [Cross Ref](#)
624. Reyes-Gibby C.C., Anderson K.O., and Todd K.H.: Risk for opioid misuse among emergency department cancer patients. *Acad Emerg Med* 2016 Feb; 23: pp. 151-158
[View In Article](#) | [Cross Ref](#)
625. Richards J.M., and Rockford R.M.: Low-dose ketamine analgesia: patient and physician experience in the ED. *Am J Emerg Med* 2013; 31: pp. 390-394
[View In Article](#) | [Cross Ref](#)
626. Richards J.R., and Rockford R.E.: Low-dose ketamine analgesia: patient and physician experience in the ED. *Am J Emerg Med* 2013; 31: pp. 390-394
[View In Article](#) | [Cross Ref](#)
627. Richards J.R., Richards I.N., Ozery G., and Derlet R.W.: Droperidol analgesia for opioid-tolerant patients. *J Emerg Med* 2011 Oct; 41: pp. 389-396
[View In Article](#) | [Cross Ref](#)
628. Richards JR, Rockford RE. Low-dose ketamine analgesia: patient and physician experience in the ED. *Am J Emerg Med.* 2013;31(2):390–4. [\[PubMed\]](#)
629. Richards JR, Rockford RE. Low-dose ketamine analgesia: patient and physician experience in the ED. *Am J Emerg Med* 2013;31:390-4.
630. Richards JR, Rockford RE. Low-dose ketamine analgesia:patient and physician experience in the ED. *Am J Emerg Med.* 2013;31(2):390–4. [\[PubMed\]](#)
631. Richter K., Adams R., Pederson D., and Bauman E.: Board 241-program innovations abstract a simulation based finale for a pain management clerkship rotation (submission #111). *Simul Healthc J Soc Simul Healthc* 2013; 8: pp. 487
[View In Article](#)
632. Rieken S, Habermehl D, Mohr A, Wuerth L, Lindel K, Weber K, Debus J, Combs SE (2011) Targeting alphanubeta3 and alphanubeta5 inhibits photon-induced hypermigration of malignant glioma cells. *Radiat Oncol* 6:132. <https://doi.org/10.1186/1748-717X-6-132>
[CrossRef](#)[PubMed](#)[PubMed Central](#)[Google Scholar](#)
633. Rieken S, Habermehl D, Wuerth L, Brons S, Mohr A, Lindel K, Weber K, Haberer T, Debus J, Combs SE (2012) Carbon ion irradiation inhibits glioma cell migration through downregulation of integrin expression. *Int J Radiat Oncol Biol Phys* 83(1):394–399. <https://doi.org/10.1016/j.ijrobp.2011.06.2004>
[CrossRef](#)[PubMed](#)[Google Scholar](#)
634. Ritcey B., Pageau P., Woo M.Y., and Perry J.J.: Regional nerve blocks for hip and femoral neck fractures in the emergency department: a systematic review. *CJEM* 2016; 18: pp. 37-47
[View In Article](#) | [Cross Ref](#)
635. Roche Laboratories Inc. Toradol oral (ketorolac tromethamine tablets). Available at:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2013/019645s019lbl.pdf. Accessed December 27, 2017.

[View In Article](#)

636. Rodgers J., Cunningham K., Fitzgerald K., and Finnerty E.: Opioid consumption following outpatient upper extremity surgery. *J Hand Surg Am* 2012 Apr; 37: pp. 645-650
[View In Article](#) | [Cross Ref](#)
637. Rodgers J., Cunningham K., Fitzgerald K., et al: Opioid consumption following outpatient upper extremity surgery. *J Hand Surg Am* 2012; 37: pp. 645-650
[View In Article](#) | [Cross Ref](#)
638. Rohatgi A. WebPlotDigitizer version 3.11. Vol 2017. Available at: <https://automeris.io/WebPlotDigitizer>. Accessed February 15, 2017.
[View In Article](#)
639. Roos H, Kellerer AM (1989) Design criteria and performance parameters of an alpha irradiation device for cell studies. *Phys Med Biol* 34(12):1823[CrossRef](#)[Google Scholar](#)
640. Rowbotham D.J., Serrano-Gomez A., and Heffernan A.: Clinical pharmacology: opioids. In Macintyre P.E. (eds): Clinical pain management (acute pain), 2nd edition. London: Hodder & Stoughton Limited, 2008. pp. 68-79
[View In Article](#)
641. Rudd R.A., Seth P.A., David F., and Scholl L.: Increases in drug and opioid-involved overdose deaths – United States, 2010–2015. *MMWR Morb Mortal Wkly Rep* 2016; 65: pp. 1445-1452
[View In Article](#) | [Cross Ref](#)
642. Rupp T, Delaney KA. Inadequate analgesia in emergency medicine. *Ann Emerg Med*. 2004; 43:494-503.
643. Rupp T, Delaney KA. Inadequate analgesia in emergency medicine. *Ann Emerg Med*. 2004; 43:494-503.
644. Rupp T., and Delaney K.A.: Inadequate analgesia in emergency medicine. *Ann Emerg Med* 2004; 43: pp. 494-503
[View In Article](#) | [Cross Ref](#)
645. Ryan H., Girion L., and S. G.: ‘You Want A Description of Hell?’ Oxycontin’s 12-hour Problem. *Los Angeles Times*, 2016 May 5.
[View In Article](#)
646. Sadove M.M., Shulman M.M., Hatano S.M., et al: Analgesic effects of ketamine administered in subdissociative doses. *Anesth Analg* 1971; 50: pp. 452-457
[View In Article](#) | [Cross Ref](#)
647. Safdar B, Choiniere M, Crandall C, et al. Impact of patient gender on pain management practices in the Emergency Department: A multi-center study. *Ann Emerg Med*. 2006;48:S121.
648. Safdar B., Degutis L., Landry K., et al: Intravenous morphine plus ketorolac is superior to either drug alone for treatment of acute renal colic. *Ann Emerg Med* 2006; 48: pp. 173-181
[View In Article](#) | [Cross Ref](#)
649. Safdar B., Degutis L.C., Landry K., et al: Intravenous morphine plus ketorolac is superior to either drug alone for treatment of acute renal colic. *Ann*

Emerg Med 2006 Aug; 48:

[View In Article](#)

650. Sato T., Matsuki A., Zsigmond E.K., and Rabito S.F.: Ketamine relaxes airway smooth muscle contracted by endothelin. Anesth Analg 1997; 84: pp. 900-906
[View In Article](#)
651. Savoia G, Coluzzi F, Di Maria C et al. Italian intersociety recommendations (SIAARTI, SIMEU, SIS 118, AISD, SIARED, SICUT, IRC) on pain management in the emergency setting. Minerva Anestesiol. 2015; 81:205-25.
652. Schmid R., Sandler A., and Katz J.: Use and efficacy of low-dose ketamine in the management of acute postoperative pain: a review of current techniques and outcomes. Pain 1999; 82: pp. 111-125
[View In Article](#) | [Cross Ref](#)
653. Schuirmann D.J.: A comparison of the two one-sided tests procedure and the power approach for assessing the equivalence of average bioavailability. J Pharmacokinet Biopharm 1987; 15: pp. 657-680
654. Schulz K.F., Altman D.G., and Moher D.: for the CONSORT Group. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. Trials 2010; 11: pp. 32
[View In Article](#)
655. Schulz K.F., Altman D.G., Moher D., and for the CONSORT Group : CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. Trials 2010; 11: pp. 32
[View In Article](#)
656. Schulz KF, Altman DG, Moher D. CONSORT 2010. Available at:
<http://www.consort-statement.org/consort-2010>. Accessed September 25, 2016.
[View In Article](#)
657. Schwartz D.R., Maroo A., Malhotra A., and Kesselman H.: Negative pressure pulmonary hemorrhage. Chest 1999; 115: pp. 1194-1197
[View In Article](#) | [Cross Ref](#)
658. Schwartz TM, Tai M, Babu KM et al. Lack of association between Press Ganey emergency department patient satisfaction scores and emergency department administration of analgesic medications. Ann Emerg Med. 2014; 64:469-81.
659. Schwartzman R., Alexander G., and Grothusen J.: The use of ketamine in complex regional pain syndrome: possible mechanisms. Expert Rev Neurother 2011; 11: pp. 719-734
[View In Article](#) | [Cross Ref](#)
660. Schwin D.A.: In Fawry G., Paul Barasch G., and Reve J.G. (eds): Cardiac pharmacology: Cardia anaesthesia, principles and practice, second edition. Philadelphia: Lippincott willkins and Williams, 2001. pp. 73-82
[View In Article](#)
661. Sehgal N., Manchikanti L., and Smith H.S.: Prescription opioid abuse in chronic pain: a review of opioid abuse predictors and strategies to curb opioid abuse. Pain Physician 2012 Jul; 15: pp. ES67-ES92
[View In Article](#)
662. Selbst SM, Clark M. Analgesics use in the emergency department. Ann Emerg Med. 1990; 19:1010-3.

663. Serinken M, Eken C, Turkcuer I et al. Intravenous paracetamol versus morphine for renal colic in the emergency department: a randomized double-blind controlled trial. *Emerg Med J*. 2012; 29:902-5.
664. Serinken M, Eken C, Turkcuer I, Elicabuk H, Uyanik E, Schultz CH. Intravenous paracetamol versus morphine for renal colic in the emergency department: a randomized double-blind controlled trial. *Emerg Med J* 2012;29:902-5.
665. Serinken M., Eken C., Turkcuer I., et al: Intravenous paracetamol versus morphine for renal colic in the emergency department: a randomised double-blind controlled trial. *Emerg Med J* 2012; 29: pp. 902-905
[View In Article](#) | [Cross Ref](#)
666. Serlin R., Mendoza T., Nakamura Y., et al: When is cancer pain mild, moderate or severe? Grading pain severity by its interference with function. *Pain* 1995; 61: pp. 277-284
[View In Article](#) | [Cross Ref](#)
667. Serlin R., Mendoza T., Nakamura Y., et al: When is cancer pain mild, moderate or severe? grading pain severity by its interference with function. *Pain* 1995; 61: pp. 277-284
[View In Article](#) | [Cross Ref](#)
668. Shams Vahdati S, Morteza Baghi HR, Ghobadi J, Rajaei Ghafouri R, Habibollahi P. Comparison of paracetamol (Apotel®) and morphine in reducing post pure head trauma headache. *Anesth Pain Med* 2014;21:e14903.
669. Sharieff GQ, Trocinski DR, Kanegaye JT, Fisher B, Harley JR. Ketamine-propofol combination sedation for fracture reduction in the pediatric emergency department. *Pediatr Emerg Care* 2007;23:881-4.
670. Sheehy KA, Lippold C, Rice AL, et al. Subanesthetic ketamine for pain management in hospitalized children, adolescents, and young adults: a single-center cohort study. *J Pain Res*. 2017;10:787-95. [\[PMC free article\]](#) [\[PubMed\]](#)
671. Sheehy KA, Lippold C, Rice AL, Nobrega R, Finkel JC, Quezado ZM. Subanesthetic ketamine for pain management in hospitalized children, adolescents, and young adults:a single-center cohort study. *J Pain Res*. 2017;10:787-95. [\[PMC free article\]](#) [\[PubMed\]](#)
672. Shimonovich S., Gigi R., Shapira A., Sarig-Meth T., Nadav D., Rozeneck M., et al: Intranasal ketamine for acute traumatic pain in the emergency department: a prospective, randomized clinical trial of efficacy and safety. *BMC Emerg Med* 2016; 16: pp. 43
[View In Article](#) | [Cross Ref](#)
673. Sin B., Tatunchak T., Paryavi M., Olivo M., et al: The use of ketamine for acute treatment of pain: a randomized, double-blind, placebo-controlled trial. *J Emerg Med* 2017; 52: pp. 601-608
[View In Article](#) | [Cross Ref](#)
674. Singer A.J., Mynster C.J., and McMahon B.J.: The effect of IM ketorolac tromethamine on bleeding time: a prospective, interventional, controlled study. *Am J Emerg Med* 2003; 21: pp. 441-443
[View In Article](#) | [Cross Ref](#)

675. Singer AJ, Mynster CJ, McMahon BJ. The effect of IM ketorolac tromethamine on bleeding time:a prospective, interventional, controlled study. Am J Emerg Med. 2003;21(5):441–3. [PubMed]
676. Smith D., Mader T., and Smithline H.: Low dose intravenous ketamine as an analgesic: a pilot study using an experimental model of acute pain. Am J Emerg Med 2001; 19: pp. 531-532
[View In Article](#) | [Cross Ref](#)
677. Smith D.J., Bouchal R.L., deSanctis C.A., Mondrode P.J., Amedro J.B., Perrotta J.M., et al: Properties of the interaction between ketamine and opiate binding sites in vivo and in vitro. Neuropharmacology 1982; 26: pp. 1253-1260
[View In Article](#) | [Cross Ref](#)
678. Smith G., Thorburn J., and Vance J.P.: The effects of ketamine on the canine coronary circulation. Anaesthesia 1979; 34: pp. 555-561
[View In Article](#) | [Cross Ref](#)
679. Smith HS. Perioperative intravenous acetaminophen and NSAIDS. Pain Med 2011;12:961–81.
680. Smith R.J., Rhodes K., Paciotti B., Kelly S., Perrone J., and Meisel Z.F.: Patient perspectives of acute pain management in the era of the opioid epidemic. Ann Emerg Med 2015; 66: pp. 246-252.e1
[View In Article](#) | [Cross Ref](#)
681. Sokoloff C., Daoust R., Paquet J., and Chauny J.M.: Is adequate pain relief and time to analgesia associated with emergency department length of stay? A retrospective study. BMJ Open 2014; 4:
[View In Article](#)
682. Soleimanipour H, Hassanzadeh K, Vaezi H et al. Effectiveness of intravenous lidocaine versus intravenous morphine for patients with renal colic in the emergency department. BMC Urol. 2012; 12:13.
683. Soleimanipour H., Hassanzadeh K., Mohammadi D.A., et al: Parenteral lidocaine for treatment of intractable renal colic: a case series. J Med Case Reports 2011 Jun 29; 5: pp. 256
[View In Article](#)
684. Soleimanipour H., Hassanzadeh K., Mohammadi D.A., et al: Parenteral lidocaine for treatment of intractable renal colic: a case series. J Med Case Rep 2011; 5: pp. 256
[View In Article](#)
685. Soleimanipour H., Hassanzadeh K., Mohammadi D.A., et al: Parenteral lidocaine for treatment of intractable renal colic: a case series. J Med Case Rep 2011; 5: pp. 256
[View In Article](#)
686. Soleimanipour H., Hassanzadeh K., Vaezi H., et al: Effectiveness of intravenous lidocaine versus intravenous morphine for patients with renal colic in the emergency department. BMC Urol 2012 May 4; 12: pp. 13
[View In Article](#)
687. Soleimanipour H., Hassanzadeh K., Vaezi H., et al: Effectiveness of intravenous lidocaine versus intravenous morphine for patients with renal colic in the emergency department. BMC Urol 2012; 12: pp. 13
[View In Article](#)
688. Soleimanipour H., Hassanzadeh K., Vaezi H., et al: Effectiveness of intravenous lidocaine versus intravenous morphine for patients with renal colic in the emergency

department. BMC Urol 2012; 12: pp. 13

[View In Article](#)

689. Soleimanpour H., Hassanzadeh K., Vaezi H., Golzari S.E., Esfanjani R.M., and Soleimanpour M.: Effectiveness of intravenous lidocaine versus intravenous morphine for patients with renal colic in the emergency department. BMC Urol 2012; 12: pp. 13

[View In Article](#)

690. Solis K.: Ethical, legal, and professional challenges posed by controlled medication seekers to healthcare providers, part 2. Am J Clin Med 2010; 7: pp. 86-92

[View In Article](#)

691. Solomon D.H., Rassen J.A., Glynn R.J., et al: The comparative safety of analgesics in older adults with arthritis. Arch Intern Med Dec 2010; 170:

[View In Article](#)

692. Somers LJ, Beckett MW, Sedgwick PM, Hulbert DC. Improving the delivery of analgesia to children in pain. Emerg Med J. 2001;18:159–161. [PMC free article] [PubMed]

693. Sorelle R. EM groups persuade Joint Commission to temper pharmacy review policy. Emerg Med News. 2007; 29:1-4. Editorial.

694. Staquet M.J.: A double-blind study with placebo control of intramuscular ketorolac tromethamine in the treatment of cancer pain. J Clin Pharmacol 1989; 29: pp. 1031-1036

[View In Article](#) | [Cross Ref](#)

695. Staquet M.J.: A double-blind study with placebo control of intramuscular ketorolac tromethamine in the treatment of cancer pain. J Clin Pharmacol 1989; 29: pp. 1031-1036

[View In Article](#) | [Cross Ref](#)

696. Staquet MJ. A double-blind study with placebo control of intramuscular ketorolac tromethamine in the treatment of cancer pain. J Clin Pharmacol. 1989;29(11):1031–6. [PubMed]

697. Stegman G.F.: Hypoxemia and suspected pulmonary edema in a Dorper ewe after diazepam-ketamine induction of anesthesia. J S Afr Vet Assoc 2000; 71: pp. 64-65

[View In Article](#)

698. Stevens D.L., King D., Laponis R., et al: Medical students retain pain assessment and management skills long after an experiential curriculum: a controlled study. Pain 2009; 145: pp. 319-324

[View In Article](#) | [Cross Ref](#)

699. Stone M.B., Price D.D., and Wang R.: Ultrasound-guided supraclavicular block for the treatment of upper extremity fractures, dislocations, and abscesses in the ED. Am J Emerg Med 2007; 25: pp. 472-475

[View In Article](#) | [Cross Ref](#)

700. Strayer 2008.

[View In Article](#)

701. Strayer R.J., and Nelson L.S.: Adverse events associated with ketamine for procedural sedation in adults. Am J Emerg Med 2008; 26: pp. 985-1028

[View In Article](#) | [Cross Ref](#)

702. Strayer R.J., Motov S.M., and Nelson L.S.: Something for pain: responsible opioid use in emergency medicine. *Am J Emerg Med* 2017; 35: pp. 337-341
[View In Article](#) | [Cross Ref](#)
703. Strigo I., Duncan G., Bushnell C., et al: The effects of racemic ketamine on painful stimulation of skin and viscera in human subjects. *Pain* 2005; 113: pp. 255-264
[View In Article](#) | [Cross Ref](#)
704. Sun Y., Li T., Wang N., et al: Perioperative systemic lidocaine for postoperative analgesia and recovery after abdominal surgery: a meta-analysis of randomized controlled trials. *Dis Colon Rectum* 2012; 55: pp. 1183-1194
[View In Article](#) | [Cross Ref](#)
705. Sutter M.E., Wintemute G.J., Clarke S.O., et al: The changing use of intravenous opioids in an emergency department. *West J Emerg Med* 2015; 16: pp. 1079-1083
[View In Article](#) | [Cross Ref](#)
706. Swing S.R., Beeson M.S., Carraccio C., et al: Educational milestone development in the first 7 specialties to enter the next accreditation system. *J Grad Med Educ* 2013; 5: pp. 98-106
[View In Article](#) | [Cross Ref](#)
707. Talati J., Tiselius H.-G., Albala D.M., et al: Urolithiasis: basic science and clinical practice. Springer Science & Business Media, Dec 22, 2012.
[View In Article](#)
708. Talwic QA. A review of the use of ketamine in pain management. *J Opioid Manag* 2013;9:379-88.
709. Tamayo-Sarver JH, Dawson NV, Hinze SW, et al. The effect of race/ ethnicity and desirable social characteristics on physicians' decisions to prescribe opioid analgesics. *Acad Emerg Med*. 2003;10:1239-1248. [PubMed]
710. Tanen D.A., Shimada M., Danish D.C., et al: Intravenous lidocaine for the emergency department treatment of acute radicular low back pain: a randomized controlled trial. *J Emerg Med* 2014; 47: pp. 119-124
[View In Article](#) | [Cross Ref](#)
711. Tanen D.A., Shimada M., Danish D.C., et al: Intravenous lidocaine for the emergency department treatment of acute radicular low back pain, a randomized controlled trial. *J Emerg Med* 2014; 47: pp. 119-124
[View In Article](#) | [Cross Ref](#)
712. Tarnow J., and Hess W.: Pulmonary hypertension and pulmonary edema caused by intravenous ketamine. *Anesthesist* 1978; 27: pp. 486-487
[View In Article](#)
713. Tawfic QA, Eipe N, Penning J. Ultra-low-dose ketamine infusion for ischemic limb pain. *Can J Anaesth*. 2014;61(1):86-7. [PubMed]
714. Tawfic QA, Eipe N, Penning J. Ultra-low-dose ketamine infusion for ischemic limb pain. *Can J Anaesth*. 2014;61(1):86-7. [PubMed]
715. Terrell K.M., Hui S.L., Castelluccio P., et al: Analgesic prescribing for patients who are discharged from an emergency department. *Pain Med* 2010 Jul; 11: pp. 1072-1077
[View In Article](#) | [Cross Ref](#)
716. Thomas S.H.: Management of pain in the emergency department. *ISRN Emerg Med* 2013; 2013:
[View In Article](#)

717. Thompson CA. Persuasive length-of-stay argument made by emergency physicians, nurses. *Am J Health-Syst Pharm*. 2007; 64:576,578. News.
718. Thomson P.: Lessons of pain: a symptom stretched across all specialties. *New Physician* 2006; 55: pp. 11-13
[View In Article](#)
719. Tintinalli J.E., Stapczynski J.S., Ma O., et al: Tintinalli's Emergency Medicine Manual. New York, NY: McGraw-Hill Medical, 2012.
[View In Article](#)
720. Tobias J.D., and Leder M.: Procedural sedation: A review of sedative agents, monitoring, and managemetn of complications. *Saudi J Anesth* 2011; 5: pp. 395-410
[View In Article](#) | [Cross Ref](#)
721. Todd K.H., Lee T., and Hoffman J.R.: The effect of ethnicity on physician estimates of pain severity in patients with isolated extremity trauma. *JAMA* 1994; 271: pp. 925-928
[View In Article](#) | [Cross Ref](#)
722. Todd K.H., Samaroo N., and Hoffman J.R.: Ethnicity as a risk factor for inadequate emergency department analgesia. *JAMA* 1993; 269: pp. 1537-1539
[View In Article](#) | [Cross Ref](#)
723. Todd KH, Deaton C, D'Adamo AP, et al. Ethnicity and analgesic practice. *Ann Emerg Med*. 2000;35:77–79. [PubMed]
724. Todd KH, Ducharme J, Choiniere M, et al. Pain in the emergency department: results of the pain and emergency medicine initiative (PEMI) multicenter study. *J Pain*. 2007; 8:460–6.
725. Todd KH, Ducharme J, Choiniere M, et al. PEMI Study Group. Pain in the emergency department: results of the pain and emergency medicine initiative (PEMI) multicenter study. *J Pain*. 2007;8:460–466. [PubMed]
726. Todd KH, Lee T, Hoffman JR. The effect of ethnicity on physician estimates of pain severity in patients with isolated extremity trauma. *JAMA*. 1994;271:925–928. [PubMed]
727. Todd KH, Samaroo N, Hoffman JR. Ethnicity as a risk factor for inadequate emergency department analgesia. *JAMA*. 1993;269:1537–1539. [PubMed]
728. Tremont-Lukats I.W., Challapalli V., McNicol E.D., et al: Systemic administration of local anesthetics to relieve neuropathic pain: a systematic review and meta-analysis. *Anesth Analg* 2005; 101: pp. 1738-1749
[View In Article](#) | [Cross Ref](#)
729. Tsui J.I., Herman D.S., Kettavong M., Alford D., Anderson B.J., and Stein M.D.: Physician introduction to opioids for pain among patients with opioid dependence and depressive symptoms. *J Subst Abuse Treat* 2010 Dec; 39: pp. 378-383
[View In Article](#) | [Cross Ref](#)
730. Turkcuer I, Serinken M, Eken C, et al. Intravenous paracetamol versus dextketoprofen in acute migraine attack in the emergency department: a randomized clinical trial. *Emerg Med J* 2014;31:182–5.
731. Turkcuer I., Serinken M., Eken C., et al: Intravenous paracetamol versus dextketoprofen in acute migraine attack in the emergency department: a randomised clinical trial. *Emerg Med J* 2014; 31: pp. 182-185
[View In Article](#) | [Cross Ref](#)

732. Turkcuer I., Serinken M., Eken C., et al: Intravenous paracetamol versus dexketoprofen in acute migraine attack in the emergency department: a randomised clinical trial. *Emerg Med J* 2014; 31: pp. 182-185
[View In Article](#) | [Cross Ref](#)
733. Turturro M.A., Paris P.M., and Seaberg D.C.: Intramuscular ketorolac versus oral ibuprofen in acute musculoskeletal pain. *Ann Emerg Med* 1995; 26: pp. 117-120
[View In Article](#) | [Cross Ref](#)
734. Turturro M.A., Paris P.M., and Seaberg D.C.: Intramuscular ketorolac versus oral ibuprofen in acute musculoskeletal pain. *Ann Emerg Med* 1995; 26: pp. 117-120
[View In Article](#) | [Cross Ref](#)
735. Tziavrangos E, Schug SA. Clinical pharmacology: other adjuvants. In: Macintyre P, et al., editors. *Clinical Pain Management (Acute Pain)* Second Edition. Hodder & Stoughton Limited; 2008. pp. 98–101.
736. Tziavrangos E, Schug SA. Clinical Pharmacology:other adjuvants. In: Macintyre P, et al., editors. *Clinical Pain Management (Acute Pain)* Second Edition. 2008 Hodder & Stoughton Limited; pp. 98–101.
737. Tziavrangos E., and Schug S.A.: Clinical pharmacology: other adjuvants. In Macintyre P. (eds): *Clinical pain management (acute pain)*, Second Edition. Hodder & Stoughton Limited, 2008. pp. 98-101
[View In Article](#)
738. Tziavrangos E., and Schug S.A.: Clinical pharmacology: other adjuvants. In Macintyre P.E., Walker S.M., and Rowbotham D.J. (eds): *Clinical pain management (acute pain)*, 2nd edition. London: Hodder & Stoughton Limited, 2008. pp. 97-98
[View In Article](#)
739. U.S. Drug Enforcement Administration website. Substance Abuse and Mental Health Services Administration. The TEDS report: substance abuse treatment admissions involving abuse of pain relievers: 1998 and 2008. Available at:
<http://www.dea.gov/divisions/dal/2010/dallas071610.pdf>. Accessed November 28, 2015.
[View In Article](#)
740. U.S. Food and Drug Administration (FDA) : FDA drug safety communication: prescription acetaminophen products to be limited to 325 mg per dosage unit; boxed warning will highlight potential for severe liver failure.
<https://www.fda.gov/Drugs/DrugSafety/ucm239821.htm>
[View In Article](#)
741. U.S. Food and Drug Administration website. Outpatient prescription opioid utilization in the U.S., years 2000-2009. Available at:
<http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/AnestheticAndLifeSupportDrugsAdvisoryCommittee/UCM220950.pdf>. Accessed November 28, 2015.
[View In Article](#)
742. Upadhyay D, Baber A, Foy M. Ketamine infusion for sickle cell pain crisis refractory to opioids:a case report and review of literature. *Ann Hematol.* 2014;93(5):769–71.
[PubMed]
743. Upadhyay D, Baber A, Foy M. Ketamine infusion for sickle cell pain crisis refractory to opioids: a case report and review of literature. *Ann Hematol.* 2014;93(5):769–71.
[PubMed]

744. Upshur C.C., Luckmann R.S., and Savageau J.A.: Primary care provider concerns about management of chronic pain in community clinic populations. *J Gen Intern Med* 2006; 21: pp. 652-655
[View In Article](#) | [Cross Ref](#)
745. Urban M., Ya Deau J., Wukovits B., et al: Ketamine as an adjunct to postoperative pain management in opioid tolerant patients after spinal fusions: a prospective randomized trial. *HSS J* 2007; 4: pp. 62-65
[View In Article](#)
746. Ury W.A., Rahn M., Tolentino V., et al: Can a pain management and palliative care curriculum improve the opioid prescribing practices of medical residents? *J Gen Intern Med* 2002; 17: pp. 625-631
[View In Article](#) | [Cross Ref](#)
747. Vahidi E., Shakoor D., Aghaie Meybodi M., and Saeedi M.: Comparison of intravenous lidocaine versus morphine in alleviating pain in patients with critical limb ischaemia. *Emerg Med J* 2015 Jul; 32: pp. 516-519
[View In Article](#) | [Cross Ref](#)
748. Vahidi E., Shakoor D., Aghaie Meybodi M., et al: Comparison of intravenous lidocaine versus morphine in alleviating pain in patients with critical limb ischaemia. *Emerg Med J* 2015; 32: pp. 516-519
[View In Article](#) | [Cross Ref](#)
749. van der Wal S.E., van den Heuvel S.A., Radema S.A., et al: The in vitro mechanisms and in vivo efficacy of intravenous lidocaine on the neuroinflammatory response in acute and chronic pain. *Eur J Pain* 2016; 20: pp. 655-674
[View In Article](#) | [Cross Ref](#)
750. Van Zee A.: The promotion and marketing of oxycontin: commercial triumph, public health tragedy. *Am J Public Health* 2009 Feb; 99: pp. 221-227
[View In Article](#) | [Cross Ref](#)
751. Vardy 2008.
[View In Article](#)
752. Vardy J.M., Dignon N., Mukherjee N., Sami D.M., Balachandran G., and Taylor S.: Audit of the safety and effectiveness of ketamine for procedural sedation in the emergency department. *Emerg Med J*. 2008; 25: pp. 579-582
[View In Article](#) | [Cross Ref](#)
753. Ventham N.T., Kennedy E.D., Brady R.R., et al: Efficacy of intravenous lidocaine for postoperative analgesia following laparoscopic surgery: a meta-analysis. *World J Surg* 2015; 39: pp. 2220-2234
[View In Article](#) | [Cross Ref](#)
754. Vigneault L., Turgeon A.F., Cote D., et al: Perioperative intravenous lidocaine infusion for postoperative pain control: a meta-analysis of randomized controlled trials. *Can J Anaesth* 2011; 58: pp. 22-37
[View In Article](#) | [Cross Ref](#)
755. Viscusi ER, Singla N, Gonzalez A, et al. IV acetaminophen improves pain management and reduces opioid requirements in surgical patients. Available at: http://www.anesthesiologynews.com/download/SR122_WM.pdf. Accessed February 6, 2016.
[View In Article](#)

756. Visvanathan T., Kluger M., Webb R., and Westhorpe R.: Crisis management during anesthesia: laryngospasm. Qual Safe Health Care 2005; 14: pp. 1136
[View In Article](#)
757. Volkow ND, McLellan TA, Cotto JH et al. Characteristics of opioid prescriptions in 2009. JAMA. 2011; 305:1299-301.
758. von Essen CF (1991) Radiation enhancement of metastasis: a review. Clin Exp Metastasis 9(2):77–104[CrossRef](#)[Google Scholar](#)
759. Vowles K.E., McEntee M.L., Julnes P.S., Frohe T., Ney J.P., and van der Goes D.N.: Rates of opioid misuse, abuse, and addiction in chronic pain: a systematic review and data synthesis. Pain 2015 Apr; 156: pp. 569-576
[View In Article](#) | [Cross Ref](#)
760. Wakai A., O'Sullivan R., and McCabe A.: Intra-articular lignocaine versus intravenous analgesia with or without sedation for manual reduction of acute anterior shoulder dislocation in adults. Cochrane Database Syst Rev 2011; undefined:
[View In Article](#)
761. Wang B, Fang Y, Jin M. Combining type-III analyses from multiple imputations. Available at: <http://support.sas.com/resources/papers/proceedings14/1543-2014.pdf>. Accessed October 28, 2016.
[View In Article](#)
762. Ward 2003.
[View In Article](#)
763. Ward J., and Standage C.: Angina pain precipitated by a continuous subcutaneous infusion of ketamine. J Pain Sympt Manage 2003; 25: pp. 6-7
[View In Article](#) | [Cross Ref](#)
764. Warner M., Chen L.H., Makuc D.M., Anderson R.N., and Miniño A.M.: Drug poisoning deaths in the United States, 1980–2008. NCHS Data Brief 2011; undefined: pp. 1-8
[View In Article](#)
765. Warner M., Hedegaard H., and Chen L.H.: Trends in drug-poisoning deaths involving opioid analgesics and heroin: United States, 1999–2012. NCHS Health E-Stat. Hyattsville (MD): National Centers for Health Statistics, 2014.
[View In Article](#)
766. Washington State Chapter of American College of Emergency Physicians. Washington emergency department opioid prescribing guidelines. www.washingtonacep.org/Postings/edopioid_abuseguidelinesfinal.pdf (accessed 2014 Nov 1).
767. Wasiak J., Mahar P.D., McGuinness S.K., et al: Intravenous lidocaine for the treatment of background or procedural burn pain. Cochrane Database Syst Rev 2014; undefined:
[View In Article](#)
768. Waxman K., Shoemaker W.C., and Lippmann M.: Cardiovascular effects of anesthetic induction with ketamine. Anesth Analg 1980; 59: pp. 355-358
[View In Article](#) | [Cross Ref](#)
769. Weant KA, Sterling E, Winstead PS et al. Establishing a pharmacy presence in the ED. Am J Emerg Med. 2006; 24:514-5. Letter.
770. Webster L.R., and Webster R.M.: Predicting aberrant behaviors in opioid-treated patients: preliminary validation of the Opioid Risk Tool. Pain Med 2005 Nov–Dec; 6: pp.

[View In Article](#) | [Cross Ref](#)

771. Weibel S., Jokinen J., Pace N.L., et al: Efficacy and safety of intravenous lidocaine for postoperative analgesia and recovery after surgery: a systematic review with trial sequential analysis. *Br J Anaesth* 2016; 116: pp. 770-783
[View In Article](#) | [Cross Ref](#)
772. Weinbroum A.A.: A single small dose of postoperative ketamine provides rapid and sustained improvement in morphine analgesia in the presence of morphine-resistant pain. *Anesth Analg* 2003; 96: pp. 789-795
[View In Article](#) | [Cross Ref](#)
773. Weiner S.G., Griggs C.A., Mitchell P.M., et al: Clinician impression versus prescription drug monitoring program criteria in the assessment of drug-seeking behavior in the emergency department. *Ann Emerg Med* 2013; 62: pp. 281-289
[View In Article](#) | [Cross Ref](#)
774. Weiner S.G., Griggs C.A., Mitchell P.M., et al: Clinician impression versus prescription drug monitoring program criteria in the assessment of drug-seeking behavior in the emergency department. *Ann Emerg Med* 2013; 62: pp. 281-289
[View In Article](#) | [Cross Ref](#)
775. Weinstein SM, Laux LF, Thornby JI, et al. Physicians' attitudes toward pain and the use of opioid analgesics: results of a survey from the Texas Cancer Pain Initiative. *South Med J*. 2000;93:479–487. [PubMed]
776. Wells G., Shea B., O'Connell D., et al: [object Object]. Ottawa, Canada: Ottawa Health Research Institute, 2004.
[View In Article](#)
777. Werner C., Reeker W., Engelhard K., Lu H., and Kochs E.: Ketamine racemate and S-(+)-ketamine: cerebrovascular effects and neuroprotection following focal ischemia. *Anesthesist* 1997; 46: pp. 55-60
[View In Article](#)
778. WGBH Educational Foundation-Getting Result Project. Getting Results. A Professional Development Course for Community College Educators. Available at: <http://www.league.org/gettingresults/web/>. Accessed Feb 20, 2011.
779. White 1982.
[View In Article](#)
780. White P.F., Way W.L., and Trevor A.J.: Ketamine-Its pharmacology and therapeutic uses. *Anesthesiology* 1982; 56: pp. 119-136
[View In Article](#) | [Cross Ref](#)
781. Wiffen P.: Clinical Pharmacology: paracetamol and compound analgesics. In Macintyre P.E., Walker S.M., and Rowbotham D.J. (eds): Clinical pain management (acute pain), 2nd edition. London: Hodder & Stoughton Limited, 2008. pp. 84-93
[View In Article](#)
782. Wightman R., Perrone J., Portelli I., and Nelson L.: Likeability and abuse liability of commonly prescribed opioids. *J Med Toxicol* 2012; 8: pp. 335-340
[View In Article](#) | [Cross Ref](#)
783. Wightman R., Perrone J., Portelli I., and Nelson L.: Likeability and abuse liability of commonly prescribed opioids. *J Med Toxicol* 2012 Dec; 8: pp. 335-340
[View In Article](#) | [Cross Ref](#)

784. Wilkerson R.G., Kim H.K., Windsor T.A., and Mareiniss D.P.: The opioid epidemic in the United States. *Emerg Med Clin North Am* 2016 May; 34: pp. e1-e23
[View In Article](#) | [Cross Ref](#)
785. Wilson J.A., Kendall J.M., and Cornelius P.: Intranasal diamorphine for paediatric analgesia assessment of safety and efficacy. *J Accid Emerg Med* 1997; 14: pp. 70-72
[View In Article](#) | [Cross Ref](#)
786. Wilson JE, Pendleton JM. Oligoanalgesia in the emergency department. *Am J Emerg Med.* 1989;7:620–623. [\[PubMed\]](#)
787. Wong C., Lau E., Palozzi L., and Campbell F.: Pain management in children: part 2 - a transition from codeine to morphine for moderate to severe pain in children. *Can Pharm J (Ott)* 2012 Nov; 145: pp. 276-279.e1
[View In Article](#) | [Cross Ref](#)
788. Wong C.S., and Wong S.H.: A new look at trigger point injections. *Anesthesiol Res Pract* 2012; 2012:
[View In Article](#)
789. Wright J.M., Price S.D., and Watson W.A.: NSAID use and efficacy in the emergency department: single doses of oral ibuprofen versus intramuscular ketorolac. *Ann Pharmacother* 1994; 28: pp. 309-312
[View In Article](#) | [Cross Ref](#)
790. Wright J.M., Price S.D., and Watson W.A.: NSAID use and efficacy in the emergency department: single doses of oral ibuprofen versus intramuscular ketorolac. *Ann Pharmacother* 1994; 28: pp. 309-312
[View In Article](#) | [Cross Ref](#)
791. Wymore ES, Casanova TJ, Broekemeier RL et al. Clinical pharmacist's daily role in the emergency department of a community hospital. *Am J Health-Syst Pharm.* 2008; 65:395-9.
792. Xia S., Chew E., Choe D., et al: No correlation between body size and hydromorphone analgesia in obese patients in ED. *Am J Emerg Med* 2015; 33: pp. 1522-1523
[View In Article](#) | [Cross Ref](#)
793. Yanni L.M., McKinney-Ketchum J.L., Harrington S.B., et al: Preparation, confidence, and attitudes about chronic noncancer pain in graduate medical education. *J Grad Med Educ* 2010; 2: pp. 260-268
[View In Article](#) | [Cross Ref](#)
794. Yeaman F, Meek R, Egerton-Warburton D, Rosengarten P, Graudins A. Sub-dissociative dose intranasal ketamine for moderate-severe pain in adult emergency department patients. *Emerg Med Australas* 2014;26:237-42.
795. Yeaman F, Oakley E, Meek R, Graudins A. Sub-dissociative dose intranasal ketamine for limb injury pain in children in the emergency department: a pilot study. *Emerg Med Australas* 2013;25:161-7.
796. Yeaman F., Oakley E., Meek R., et al: Sub-dissociative dose intranasal ketamine for limb injury pain in children in the emergency department:a pilot study. *Emerg Med Australas* 2013; 25: pp. 161-167
[View In Article](#) | [Cross Ref](#)

797. Yeh Y.C., and Reddy P.: Clinical and economic evidence for intravenous acetaminophen. *Pharmacotherapy* 2012; 32: pp. 559-579
[View In Article](#) | [Cross Ref](#)
798. Yeh Y.C., and Reddy P.: Clinical and economic evidence for intravenous acetaminophen. *Pharmacotherapy* 2012; 32: pp. 559-579
[View In Article](#) | [Cross Ref](#)
799. Yeh YC, Reddy P. Clinical and economic evidence for intravenous acetaminophen. *Pharmacotherapy* 2012;32:559-79.
800. Yen K, Kim M, Stremski ES, Gorelick MH. Effect of ethnicity and race on the use of pain medications in children with long bone fractures in the emergency department. *Ann Emerg Med*. 2003;42:41–47. [PubMed]
801. Yong C., Hongguang N., Hong M., Chen L., Zhang L., Wang J., et al: Ketamine inhibits lung fluid clearance through reducing alveolar sodium transport. *J Biomed Biotech* 2011; 2011: pp. 1-6
[View In Article](#)
802. Younge P.A., Nicol M.F., Kendall J.M., and Harrington A.P.: A prospective randomized pilot comparison of intranasal fentanyl and intramuscular morphine for analgesia in children presenting to the emergency department with clinical features. *Emerg Med (Fremantle)* 1999; 11: pp. 90-94
[View In Article](#) | [Cross Ref](#)
803. Zacny J.P., and Lichtor S.A.: Within-subject comparison of the psychopharmacological profiles of oral oxycodone and oral morphine in non-drug-abusing volunteers. *Psychopharmacology (Berl)* 2008; 196: pp. 105-116
[View In Article](#) | [Cross Ref](#)
804. Zacny J.P., and Lichtor S.A.: Within-subject comparison of the psychopharmacological profiles of oral oxycodone and oral morphine in non-drug-abusing volunteers. *Psychopharmacology (Berl)* 2008 Jan; 196: pp. 105-116
[View In Article](#) | [Cross Ref](#)
805. Zare MA, Ghalyae AH, Fathi M, Farsi D, Abbasi S, Hafezimoghadam P. Oral oxycodone plus intravenous acetaminophen versus intravenous morphine sulfate in acute bone fracture pain control: a double-blind placebo-controlled randomized clinical trial. *Eur J Orthop Surg Traumatol* 2014;24:1305-9.
806. Zed PJ, Filiault L. Clinical outcomes and patient satisfaction of a pharmacist managed, emergency department-based outpatient treatment program for venous thromboembolic disease. *CJEM*. 2008; 10:10-7.
807. Zempsky W.T., Loiselle K.A., Corsi J.M., and Hagstrom J.N.: Use of low-dose ketamine infusion for pediatric patients with sickle cell disease-related pain. *Clin J Pain* 2010; 26: pp. 164-167
[View In Article](#)
808. Zempsky WT, Loiselle KA, Corsi JM, et al. Use of low-dose ketamine infusion for pediatric patients with sickle cell disease-related pain: a case series. *Clin J Pain*. 2010;26(2):163–7. [PubMed]
809. Zempsky WT, Loiselle KA, Corsi JM, Hagstrom JN. Use of low-dose ketamine infusion for pediatric patients with sickle cell disease-related pain:a case series. *Clin J Pain*. 2010;26(2):163–7. [PubMed]

810. Zielmann S., Kazmaier S., Schnüll S., and Weyland A.: S-(+)-ketamine and circulation. *Anaesthesia* 1997; 46: pp. 43-46
[View In Article](#) | [Cross Ref](#)
811. Zorzela L., Loke Y.K., Ioannidis J.P., et al: PRISMA harms checklist: improving harms reporting in systematic reviews. *BMJ* 2016; 352: pp. i157
[View In Article](#) | [Cross Ref](#)